

27th International Symposium on Analytical and Environmental Problems



PROCEEDINGS OF THE

27th International Symposium
on Analytical and Environmental Problems

Szeged, Hungary
November 22-23, 2021



University of Szeged

Edited by:
Tünde Alapi
Róbert Berkecz
István Ilisz

Publisher:
University of Szeged, H-6720 Szeged, Dugonics tér 13,
Hungary

ISBN 978-963-306-835-9

2021.
Szeged, Hungary

***The 27th International Symposium on Analytical and
Environmental Problems***

Organized by:

SZAB Kémiai Szakbizottság Analitikai és Környezetvédelmi Munkabizottsága

Supporting Organizations

*Institute of Pharmaceutical Analysis, University of Szeged
Department of Inorganic and Analytical Chemistry, University of Szeged*

Symposium Chairman:

István Ilisz, DSc

Honorary Chairman:

Zoltán Galbács, PhD

Organizing Committee:

István Ilisz, DSc

professor of chemistry

University of Szeged, Institute of Pharmaceutical Analysis

Tünde Alapi, PhD

assistant professor

University of Szeged, Department of Inorganic and Analytical Chemistry

Róbert Berkecz, PhD

assistant professor

University of Szeged, Institute of Pharmaceutical Analysis

Scientific Committee:

István Ilisz, DSc

Tünde Alapi, PhD

Róbert Berkecz, PhD

Daniela Sojic Merkulov, PhD

associate professor

*University of Novi Sad, Faculty of Sciences, Department of Chemistry, Biochemistry and
Environmental Protection*

Lecture Proceedings

COMBINED EXPERIMENTAL AND DFT STUDY OF LITHIUM-INDIUM-OXIDE STRUCTURE AND VIBRATIONAL PROPERTIES

Robert Vigi¹, Ljubica Đaćanin Far², Svetlana Lukić-Petrović¹, and Tamara Ivetić¹

¹University of Novi Sad, Faculty of Sciences, Department of Physics, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia

²University of Belgrade, Vinča Institute of Nuclear Sciences, Center of Excellence for Photoconversion, P.O. Box 522, 11000 Belgrade, Serbia
e-mail: robert1995@uns.ac.rs

Abstract

A promising lithium-indium-oxide (LiInO₂) wide band-gap semiconductor for scintillating detection, photoluminescence, and photocatalysis [1-3] was prepared by a mechanochemical solid-state synthetic procedure that can be found elsewhere [3]. Its structure and morphology were investigated by using X-ray diffraction (XRD), scanning electron microscopy (SEM), and Raman spectroscopy. SEM images show agglomerates of relatively uniform size of around 300 nm spherical-shaped particles of LiInO₂ powder, while the XRD pattern confirmed the formation of the nanocrystalline tetragonal structure with $I4_1/amd$ space group (no. 141) symmetry. Detailed vibration analysis, together with the assignments of the band modes, was performed through the best-fit match of the experimental and density functional theory (DFT) calculated Raman spectrum. Geometry optimizations and vibrational frequencies calculations were conducted using B97-1 functional correlation [4] and LanL2DZ was used as a basis set.

Acknowledgements

The authors acknowledge the financial support of the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant No. 451-03-9/2021-14/ 200125).

References

- [1] Lj.R. Đaćanin, M.D. Dramićanin, M.G. Nikolić, M. Mitrić, D.M. Petrović, S.R. Lukić, Phys. Status Solidi (C) Curr. Top. in Solid State Phys. 8 (2011) 2830.
- [2] Lj. Đaćanin, S.R. Lukić-Petrović, D.M. Petrović, M.D. Dramićanin, Phys. Scripta 85 (2012) 065703.
- [3] Lj.R. Đaćanin Far, N.L. Finčur, T.B. Ivetić, B.F. Abramović, D.Štrbac, O.Bosak, S.R. Lukić-Petrović, Rom. J. Phys. 65 (2020) 601.
- [4] Hamprecht, F. A., Cohen, A. J., Tozer, D. J., Handy, N. C. J. Chem. Phys. 1998,109, 6264.