

ANTIOXIDANT STATUS IN THE BLOOD OF PSYCHOSOCIALLY STRESSED RATS TREATED WITH HONEY

Ljubica GAVRILOVIĆ^{1*}, Vesna STOJILJKOVIĆ¹, Vojislav STANIĆ², Dragoljub JOVANOVIĆ³, Snežana PEJIĆ¹, Branka BOROVIĆ⁴, Snežana B. PAJOVIĆ¹

¹University of Belgrade, “Vinča” Institute of Nuclear Sciences, National Institute of the Republic of Serbia, Department of Molecular Biology and Endocrinology, Belgrade, Serbia; ²University of Belgrade, Institute of Nuclear Sciences “Vinča”, National Institute of the Republic of Serbia, Department of Radiation and Environmental Protection, Belgrade, Serbia; ³University of Belgrade, Faculty of Veterinary Medicine, Belgrade, Serbia; ⁴Institute of Meat Hygiene and Technology, Belgrade, Serbia

(Received 01 August, Accepted 17 November 2023)

Linden honey represents a unique honey variety valued for its nutritional benefits, distinctive taste and aroma. Phenols, polyphenols, flavonoids, ascorbic acid and phenolic acids in honey have antioxidant activities. This study aimed to investigate the effects of linden honey on the activities of antioxidant enzymes catalase (CAT) and glutathione peroxidase (GPx), as well as on the concentration of malondialdehyde (MDA) in individually housed animals. The investigated parameters were quantified using spectrophotometric method for determination of enzyme activities and MDA concentration in the blood. We found that treatment with linden honey in the socially isolated animals significantly increased the enzyme activities of CAT and GPx, and significantly decreased the concentration of MDA. The modulation of CAT and GPx activities in socially isolated animals treated with linden honey may be very important for understanding the role of honey in the capacity of antioxidant defense system to increase and maintain its stability in psychosocial stress conditions. Our results may be important in biomedical research for understanding the role of honey in the amelioration of oxidative stress.

Keywords: honey, blood, social isolation, enzymes activity, antioxidant status

INTRODUCTION

Social isolation produced by rats' individual housing represents a mild psychosocial stress and resembles perceived isolation observed in depressive disorders [1]. Many studies have shown that social isolation and environmental changes are risk factors in human depression [2]. In addition, COVID-19 pandemic that resulted in quarantine and social isolation has led to the rise in anxiety [3] and also to an increase in posttraumatic

*Corresponding author: e-mail: gljubica@vin.bg.ac.rs

stress disorder [4]. Chronic social isolation in adult male Wistar rats causes depressive-like and anxiety-like behaviors [5]. It is known that psychosocial stress is associated with increased oxidative stress and decreased antioxidant capacity [6]. For example, the increase in hepatic reactive oxygen species (ROS) amount in individually housed rats might be due to social isolation-induced dysfunctions of antioxidant enzymes [7]. Additionally, isolated rats showed a significant increase in malondialdehyde (MDA) [8] which is the end product of lipid peroxidation and biomarker of free radicals. It is known that long-term isolation induces changes in the activities of endogenous antioxidants. For example, the mentioned stressor changes hippocampal copper, zinc-superoxide dismutase (CuZnSOD) and catalase (CAT) activity [9], as well as hepatic glutathione-dependent defense system [10]. However, very little is known about the ability of natural exogenous antioxidants to reduce increased oxidative stress and decreased antioxidant capacity caused by psychosocial stressors. The doses of honey used in this study have been reported in previous studies [11,12], as well as that it is considered good natural dietary source of antioxidants [13]. It was confirmed that honey contains vitamins including vitamin C as an antioxidant that can capture free radicals [14]. The literature data confirm that the identified compounds in linden honey include monoterpenes (hydrocarbons, ethers, aldehydes, acids, and bifunctional derivatives), isoprenoids, aromatic compounds (phenylpropanoids, phenols) and products degraded from fatty acids to alkaloids [15]. In addition, the results of Gašić and colleagues [16] have shown that samples of honey from Vojvodina and Zlatibor region were distinguished from the honey varieties from the rest of Serbia due to the presence of dicaffeoylquinic acid, ellagic acid, caffeic acid phenethyl ester, and chlorogenic acid. For this reason, in this study we examined the antioxidant status of the blood of psychosocially stressed rats treated with linden honey from the area of Fruška Gora. In addition, the property of linden honey to inhibit lipid peroxidation, as well as to capture hydroxyl and superoxide anion radicals was examined by the “Laboratory for testing natural resources and pharmacologically and biologically active compounds”, Novi Sad. Because of the direct involvement of oxidative stress in anxiety-like and depressive-like behavior in stress conditions, detecting the changes of activity of the antioxidant enzymes, as well as concentration of MDA in psychosocially stressed rats treated with linden honey may be very important in the research of the role of linden honey in maintaining antioxidant capacity in pathological conditions.

In this study we examined the activity of the antioxidant enzymes CAT and GPx, and concentrations of MDA in the blood of chronically stressed rats treated with linden honey.

MATERIALS AND METHODS

Animals

In the experiment adult male rats of Wistar strain, 11-week-old and weighing 300-350 g were used. The animals were maintained under standard vivarium conditions in a temperature-controlled room (22 ± 1.0 °C) and 12 h/12 h light/dark cycle, with

water and food *ad libitum*. Handling of animals was done quickly and carefully to avoid unnecessary discomfort. The rats were habituated to handling and treated as ethically as possible, according to the recommendations of the Ethical Committee of the “Vinča” Institute of Nuclear Sciences, Belgrade, Serbia, which follows the guidelines of the Serbian Society for the Use of Animals in Research and Education. The Committee has approved the planned experiment (06/2023).

Experiment design

Ten (10) animals were randomly divided into two groups. First group, individually housed group (IH group, n=5) of animals was subjected to social isolation with a single animal per cage for 28 days. The visual and olfactory communication among the isolated rats was reduced to a minimal level.

Second group, individually housed group + honey group (IH+HONEY group, n=5), consisted of individually housed animals given 1,5 ml/kg honey orally for 28 days.

The linden honey used in this experiment was collected in 2022 by dr Vojislav Stanić in an apiary located in Vojvodina-Fruška Gora region and stored at room temperature in dark until the experiment. The honey was mixed with distilled water in a ratio of 1:1. The doses of honey used in this study have been reported by Abdulmajeed et al. [11] and Halawa et al. [12].

After the described treatment, blood samples were collected from both groups. Blood was sampled from the rat's tail (into heparinized tubes). After blood collection, the were returned to their cages alive.

Determination of malondialdehyde (MDA), catalase (CAT) activity and glutathione peroxidase (GPx) activity

Determination of MDA level was performed using the spectrophotometric method previously described by Siddique et al. [17]. All sample values were read on the spectrophotometer at 586 nm. Malondialdehyde concentration was expressed as $\mu\text{mol/ml}$ of plasma.

CAT activity was determined by the method of Beutler [18]. All samples were read on the spectrophotometer at 230 nm. The final result for CAT activity was expressed as units per milligram of hemoglobin (U/mg Hb).

Determination of activity was performed using methods previously described by Agergaard and Jensen [19]. All samples were read on spectrophotometer at 366 nm. The final result for GPx activity was expressed as units per gram of hemoglobin (U/g Hb).

Data analysis

The data are presented as means \pm S.E.M. The differences in the activity of enzymes (CAT and GPx) between IH group and IH+HONEY group were analyzed by t-test.

Statistical analysis was carried out using the SigmaPlot 10.0 with SigmaStat integration. The statistical significance was accepted at $p < 0.05$.

RESULTS

The honey treatment in individually housed animals (IH+HONEY group) resulted in significantly higher CAT activity (26%) (Figure 1a), and GPx activity (31%) (Figure 1b) compared with individually housed animals (IH group). In addition, socially isolated animals treated with honey (IH+HONEY group) had a 35% lower concentration of MDA (Figure 2) compared with individually housed animals (IH group).




Figure 1. The effects of linden honey on the enzyme activities of catalase (CAT) [a] and glutathione peroxidase (GPx) [b] in the blood of individually housed animals. The values are means \pm S.E.M. of 10 rats. Statistical significance: $+p < 0.05$ IH+HONEY group *vs.* IH group (t-test). The final result for CAT activity was expressed as units per milligram of hemoglobin (U/mg Hb) and the final result for GPx activity was expressed as units per gram of hemoglobin (U/g Hb).




Figure 2. The effects of linden honey on the concentrations of malondialdehyde (MDA) in blood of individually housed animals. The values are means \pm S.E.M. of 10 rats. Statistical significance: $+p < 0.05$ IH+HONEY group *vs.* IH group (t-test). The final result for the concentration of MDA was expressed as $\mu\text{mol/ml}$ of plasma.

DISCUSSION

In our previous study we confirmed that social isolation significantly increases the concentration of MDA in the left adrenal medulla [20]. In addition, the literature data confirm that social isolation influence the occurrence of oxidative stress [21]. Also, Bove et al. [7] observed that isolated rats showed significant elevations in ROS and increased concentration of MDA in the liver, which confirms that social isolation triggers oxidative stress. It is known that the increased lipid peroxidation in isolated rats can be a consequence of the dysfunction of antioxidant enzymes, which confirms that the antioxidant defense system is insufficient to keep the levels of ROS low [7]. Several studies have confirmed that honey supplementation has therapeutic effects under pathological conditions [22,23]. In the present study we observed that in response to linden honey antioxidant enzymes underwent modulation. We found that socially isolated animals treated with linden honey showed increased CAT and GPx activity compared with individually housed animals. It is known that CAT and GPx protect tissues from highly reactive hydroxyl radical derived from H₂O₂ [24]. Our results are in accordance with the reports of Oyefuga and colleagues [25] who found that supplementation with honey significantly decreased the lipid peroxidation and increased antioxidant enzymes activity in the brain. The results presented in this study indicated that linden honey increases antioxidant activity and that it is possibly involved in the protection from oxidative stress. The literature data have shown that the effect of honey to prevent of oxidative damage might be due to its phenolic and non-phenolic antioxidant content [26]. It is known that phenols, polyphenols, flavonoids, catalase, ascorbic acid and phenolic acids in honey have antioxidant activities [27,28]. "The laboratory for testing natural resources and pharmacologically and biologically active compounds" Novi Sad confirmed that the linden honey used in our experiment has the ability to capture hydroxyl radicals $IC_{50}=1.116 \text{ mg/ml} \pm 0.157$.

The ability of honey to neutralize hydroxyl radicals was tested using the spectrophotometric method previously described by Pintać et al. [29]. Recent studies have confirmed that honey contains CAT, as well as that the antioxidant defense mechanism of honey can be attributed to CAT [30]. Also, as reported by Gheldoff and Engeseth [31], honey contains selenium. Costa-Silva et al. [32] found that increased GPx might be due to the presence of selenium in honey which is a cofactor for GPx activity. In addition, Halawa et al. [12] have suggested that honey exerts its protective role through its antioxidant mechanism and through restoring glutathione activity in rats. Namely, honey is a good natural source of sulfhydryl (SH) groups. Also, it is known that the most important endogenous antioxidant is glutathione, as well as that the SH group of glutathione is important as a direct scavenger of ROS [33]. Abdulmajeed et al. [11] found that administration of honey improves antioxidant activities as shown by increased brain SOD and glutathione-dependent defense system. An important result of this study is that honey treatment in animals exposed to social isolation significantly decreases concentration of MDA in blood. Decreased levels of

MDA in individually housed animals treated with linden honey, found in our study, confirm that honey is involved in the reduction of oxidative stress in psychosocial stress conditions. The reduced MDA levels are probably the consequence of increased CAT and GPx activity. In addition, the “Laboratory for testing natural resources and pharmacologically and biologically active compounds” Novi Sad, confirmed that the linden honey we used in our experiment has the ability to inhibit lipid peroxidation ($24.5\% \pm 3.2$ at the concentration of honey 0.179 mg/ml; spectrophotometric method previously described by Pintać *et al.* [29]). Our results confirm that linden honey increases the antioxidant activity and can protect from oxidative stress. Also, the latest results confirmed that linden honey is a source of numerous vitamins, minerals, organic and inorganic acids [34]. For example, Farkas *et al.* [34] showed that linden honey samples displayed significantly higher levels of caffeic, chlorogenic, ferulic and *p*-coumaric acids compared to acacia and goldenrod honeys. In addition, apigenin, chrysin and galangin were measured in both Hungarian and Serbian linden honey samples [34]. These results suggest that the geographical origin can strongly influence the composition of characteristic compounds which can contribute to the health benefits of honey [34].

CONCLUSIONS

In the present study we found that treatment with linden honey in socially isolated animals significantly increased the enzyme activities of CAT and GPx, and significantly decreased the concentration of MDA. The modulation of CAT and GPx activities in socially isolated animals treated with linden honey may be very important for understanding the role of honey in the capacity of the antioxidant defense system to increase and maintain its stability in psychosocial stress conditions. The understanding of the influence of honey on redox balance regulation, might contribute to the development of therapeutic strategies, aiming at reduction of oxidative stress in pathological conditions.

Acknowledgmentst

This work was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Research Grant No.451-03-47/2023-01/200017).

Authors' contributions

LjG participated in the experiment, in the statistical analysis, designed and wrote the manuscript. VS participated in the experiment, participated to draft the statistical analysis, helped to draft the manuscript. VS helped to draft the statistical analysis. DJ participated in the experiment. SP helped to draft the manuscript. BB participated in the analysis of the results. SBP helped to draft the manuscript. All authors read and approved the final manuscript.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical Statement

The Ethical Committee for the care and use of laboratory animals of the Institute of Nuclear Sciences “Vinča”, Belgrade, Serbia has approved the planned experiment. The opinion (06/2023) of the Ethical Committee is that the use of laboratory animals is ethically justified for conducting the experiment. The Ethical Committee of the “Vinča” Institute of Nuclear Sciences “Vinča”, Belgrade, Serbia, follows the guidelines of the Serbian Society for the Use of Animals in Research and Education.

REFERENCES

1. Hawkley LC, Cole SW, Capitano JP, Norman GJ, Cacioppo JT: Effects of social isolation on glucocorticoid regulation in social mammals. *Horm Behav* 2012, 62:314–323.
2. Ishida H, Mitsui K, Nukaya H, Matsumoto K, Tsuji K: Study of active substances involved in skin dysfunction induced by crowding stress. I. Effect of crowding and isolation on some physiological variables, skin function and skin blood perfusion in hairless mice. *Biol Pharm Bull* 2003, 26(2):170-181.
3. Twenge JM, Joiner TE: U.S. Census Bureau-assessed prevalence of anxiety and depressive symptoms in 2019 and during the 2020 COVID-19 pandemic. *Depress Anxiety* 2020.
4. Yuan K, Gong YM, Liu L, Sun YK, Tian SS, Wang YJ, Zhong Y, Zhang AY, Su SZ, Liu XX, Zhang YX, Lin X, Shi L, Yan W, Fazel S, Vitiello MV, Bryant RA, Zhou XY, Ran MS, Bao YP, Shi J, Lu L: Prevalence of posttraumatic stress disorder after infectious disease pandemics in the twenty-first century, including COVID-19: a meta-analysis and systematic review. *Mol Psychiatry* 2021.
5. Zlatković J, Todorović N, Bošković M, Pajović SB, Demajo M, Filipović, D: Different susceptibility of prefrontal cortex and hippocampus to oxidative stress following chronic social isolation stress. *Mol Cell Biochem* 2014, 393(1–2):43–57.
6. Shuaib DM, Khalid BEA: Effect of Social Isolation on Rat Ovary and the Possible Protective Role of Melatonin: Light, Electron Microscopic, Biochemical and Molecular Study. *Med J Cairo Univ* 2020, 88(2):529-540.
7. Bove M, Lama A, Schiavone S, Pirozzi C, Tucci P, Sikora V, Trinchese G, Corso G, Morgese M G, Trabace L: Social isolation triggers oxidative status and impairs systemic and hepatic insulin sensitivity in normoglycemic rats. *Biomed Pharmacother* 2022, 149:112820.
8. Ali A, Khalil MG, Elariny HA, Abu-Elfotuh K: Study on Social Isolation as a Risk Factor in Development of Alzheimer’s Disease in Rats. *Brain Disord Ther* 2017, 6:2.
9. Pajović SB, Pejić S, Stojilković V, Gavrilović L, Dronjak S, Kanazir DT: Alternations in hippocampal antioxidant enzyme activities and symphato-adrenomedullary system of rats in response to different stress models. *Physiol Res* 2006, 55:453-460.
10. Todorović N, Tomanović N, Gass P, Filipović D: Olanzapine modulation of hepatic oxidative stress and inflammation in socially isolated rats. *European Journal of Pharmaceutical Sciences* 2016, 1(81):94-102.

11. Abdulmajeed WI, Sulieman HB, Zubayr MO, Imam A, Amin A, Biliaminu SA, Oyewole LA, Owoyele BV: Honey prevents neurobehavioural deficit and oxidative stress induced by lead acetate exposure in male wistar rats- a preliminary study. *Metab Brain Dis* 2016, 31:37-44.
12. Halawa HM, El-Nefawy NE, Makhlof NA, Mady AA: Evaluation of honey protective effect on lead induced oxidative stress in rats. *JASMR* 2009, 4(2):197-208.
13. Gharzouli K, Gharzouli A, Amira S, Khennouf S: Protective effect of mannitol, glucose-fructose– maltose mixture, and natural honey hyperosmolar solutions against ethanol – induced gastric mucosal damage in rats. *Exp Toxic Pathol* 2001, 53:175-180.
14. Fajrilah BR, Indrayani UD, Djama Q: The Effect of Honey on Plasma Malondialdehyde (MDA) Level on Alloxan-Induced hyperglycemic Rats An Experimental studies in rats Galur Wistar White Males. *Sains medika* 2013, 5(2):98-100.
15. Naef R, Jaquier A, Velluz A, Bachofen B: From the linden flower to linden honey--volatile constituents of linden nectar, the extract of bee-stomach and ripe honey. *Chem Biodivers* 2004, 1(12):1870-1879.
16. Gašić U, Kečkeš S, Dabić D, Trifković J, Milojković-Opšenica D, Natić M, Tešić Ž: Phenolic profile and antioxidant activity of Serbian polyfloral honeys. *Food Chem* 2014, 145:599-607.
17. Siddique YH, Ara G, Afzal M: Estimation of lipid peroxidation induced by hydrogen peroxide in cultured human lymphocytes. *Dose-Response* 2012, 10:1-10.
18. Beutler E: Catalase. *Red Cell Metabolism*. 3rd ed. *A Manual of Biochemical Methods*. Orlando, Fla, USA: Grune & Stratton 1982.
19. Agergaard N, Jensen PT: Procedure for blood glutathione peroxidase determination in cattle and swine. *Acta Vet Scand* 1982, 23(4):515-527.
20. Gavrilović L, Stojiljković V, Pejić S, Spasojević Tišma V, Nikolić D, Pajović SB: Differences in the functional activity and redox homeostasis between the left and right adrenal gland of rats exposed to chronic isolation stress. *Acta Vet – Beograd* 2022, 72(2):224-234.
21. Schiavone S, Jaquet V, Trabace L, Krause KH: Severe life stress and oxidative stress in the brain: from animal models to human pathology. *Antioxid Redox Signal* 2013, 18(12):1475-1490.
22. Erejuwa OO, Sulaiman SA, Wahab MSA, Sirajudeen KNS, Salleh MSM, Gurtu S: Differential responses to blood pressure and oxidative stress in streptozotocin-induced diabetic wistar-kyoto rats and spontaneously hypertensive rats: effects of antioxidant (Honey) treatment. *Int J of Mol Sci* 2011, 12(3):1888–1907.
23. Erejuwa OO, Sulaiman SA, AbWahab MS, Sirajudeen KNS, Salleh S, Gurtu S: Honey Supplementation in Spontaneously Hypertensive Rats Elicits Antihypertensive Effect via Amelioration of Renal Oxidative Stress. *Oxid Med Cell Longev* 2012, Article ID 374037, 14 pages doi:10.1155/2012/374037
24. Haber F, Weiss J: The catalytic decomposition of hydrogen peroxide by iron salts. *P Roy Soci* 1934, 147:332–351.
25. Oyefuga OH, Ajani EO, Salau BA, Agboola F, Adebawo OO: Honey consumption and its anti-ageing potency in white Wister albino rats. *Scholarly Journal of Biological Science* 2012, 1(2):15-19.
26. El-Shafey AAM, Seliem MME, El-Zawahry SAM, Shahen EMS, Mahmoud DEM: Effect of honey bee on some antioxidant enzymes and lipid profile in hypercholesterolemic male albino rats. *ZUMJ* 2015, 21(1):68-78.

27. Bogdanov S, Jurendic T, Sieber R, Gallmann P: Honey for nutrition and health: a review. *J Am Coll Nutr* 2008, 27(6):677–689.
28. El Rabey HA, Al Seeni MN, AlSolamy SM: Bees' honey protects the liver of male rats against melamine toxicity. *BioMed Res Intern* 2013, ID 786051: 8.
29. Pintac D, Četojević-Simin D, Berežni S, Orčić D, Mimica-Dukić M, Lesjak M. Investigation of the chemical composition and biological activity of edible grapevine (*Vitis vinifera* L.) leaf varieties. *Food Chem* 2019, 268, 686-695. 10.1016/j.foodchem.2019.02.049
30. Mir B, Mushtaq A, Rizwan S, Bashir F: Comparison of Antioxidant Enzymes in Natural and Commercial Honey. *PJMLS* 2022, 5(1):31-38.
31. Gheldof N, Engeseth NJ: Antioxidant capacity of honeys from various floral sources based on the determination of oxygen radical absorbance capacity and inhibition of in vitro lipoprotein oxidation in human serum samples. *J Agric Food Chem* 2002, 50(10):3050–3055.
32. Costa-Silva F, Maia M, Matos CC, Calçada E, Barros AIRNA, Nunes FM: Selenium content of Portuguese unifloral honeys. *J Food Comp Anal* 2011, 24:351-355.
33. Schulz JB, Lindenau J, Seyfried J: Glutathione oxidative stress and neurodegeneration. *Eur J Biochem* 2000, 267:4904-4911.
34. Á Farkas, G Horváth, M Kuzma, M Mayer, M Kocsis: Phenolic compounds in Hungarian acacia, linden, milkweed and goldenrod honeys. *Curr Res Food Sci* 2023, 6:100526. doi: 10.1016/j.crfs.2023.100526

ANTIOKSIDATIVNI STATUS U KRVI PACOVA IZLAGANIH PSIHOSOCIJALNOM STRESU TRETIRANIH MEDOM

Ljubica GAVRILOVIĆ, Vesna STOJILJKOVIĆ, Vojislav STANIĆ, Dragoljub JOVANOVIĆ, Snežana PEJIĆ, Branka BOROVIĆ, Snežana B. PAJOVIĆ

Lipov med predstavlja jedinstvenu vrstu meda cenjenu zbog svoje nutritivne prednosti, specifičnog ukusa i mirisa. Fenoli, polifenoli, flavonoidi, askorbinska kiselina i fenolne kiseline u medu imaju antioksidativno dejstvo. Ovo istraživanje je imalo za cilj da ispita uticaj lipovog meda na aktivnosti antioksidativnih enzima katalaze (CAT) i glutation peroksidaze (GPx), kao i na koncentraciju malondialdehida (MDA) kod individualno smeštenih životinja. Ispitivani parametri su kvantifikovani spektrofotometrijskim metodama. Ustanovili smo da tretman lipovim medom, kod socijalno izolovanih životinja, značajno povećava aktivnost enzima CAT i GPx, ali da značajno smanjuje koncentraciju MDA. Modulacija aktivnosti CAT i GPx u krvi socijalno izolovanih životinja tretiranih lipovim medom, može biti veoma važna za razumevanje uloge meda u sposobnosti antioksidativnog odbrambenog sistema da poveća i održi njegovu stabilnost u uslovima psihosocijalnog stresa. Naši rezultati mogu biti važni u biomedicinskim istraživanjima za razumevanje uloge meda u smanjenju oksidativnog stresa.