

Article

Non-Abelian Topological Approach to Non-Locality of a Hypergraph State

Vesna Berec ^{1,2}

¹ Institute of Nuclear Sciences Vinca, P.O. Box 522, 11000 Belgrade, Serbia, E-Mail: vberec@vinca.rs; Tel.: +381-11-3408-352

² University of Belgrade, Studentski trg 1, 11000 Belgrade, Serbia; E-Mail: bervesn@gmail.com

Academic Editors: Jiannis Pachos, Demosthenes Ellinas, Giorgio Kaniadakis and Antonio M. Scarfone

Received: 16 February 2015 / Accepted: 8 May 2015 / Published: 15 May 2015

Abstract: We present a theoretical study of new families of stochastic complex information modules encoded in the hypergraph states which are defined by the fractional entropic descriptor. The essential connection between the Lyapunov exponents and d -regular hypergraph fractal set is elucidated. To further resolve the divergence in the complexity of classical and quantum representation of a hypergraph, we have investigated the notion of non-amenability and its relation to combinatorics of dynamical self-organization for the case of fractal system of free group on finite generators. The exact relation between notion of hypergraph non-locality and quantum encoding through system sets of specified non-Abelian fractal geometric structures is presented. Obtained results give important impetus towards designing of approximation algorithms for chip imprinted circuits in scalable quantum information systems.

Keywords: non-Abelian group; hypergraph state; topological system; non-locality; geometry information

1. Introduction

The field of algebraic topology has passed an essential evolution in attaching algebraic objects to topological spaces starting from the simple graph models associated to vertices and edges in terms of Laplacian matrices, leading to higher-order dimensional structures modified through simplices or cliques, finalizing with an elegant mature representation of non-local hypergraph states [1]. From the aspect of the algebraic topology, the dimension of the specific measure for the hypergraph (set system)

is a function of the corresponding metric [2]. As a natural extension former implicates that for algebraic dynamical system the entropy of the measure depends on set system evolution process upon its metric spaces [3], *i.e.*, the metric space represents the natural constituent assembly of the set system dimension in dynamical framework. Moreover, for algebraic dynamical systems the dimension directly relates to entropy via geometric concept of the Lyapunov exponents [4,5], which can be successfully applied to quantify the set system complexity in a sense of the topological [6,7] as well as the metric entropy [8].

In general sense the topological entropy represents quantitative measure of complexity for continuous map defined on compact metric spaces of dynamical system [2]. While the metric entropy quantifies the number of typical orbits, the topological entropy represents the exponential growth rate of the number of orbit segments that are distinguishable with finite resolution. Hence, dynamical property of the maps is regulated by the topological entropy, *i.e.*, between distinguishable orbits it measures growth rate of the number of different orbits of length n in infinite limit. Specifically, topological entropy as a non-causal measure of exponential increase of the number of maximal intervals of monotonicity is introduced for a piecewise interval monotone map [9,10].

Important feature of the set system topological entropy is its close relation to periodic orbits. For each periodic orbit P of continuous map f [11] given on the interval I , it can be assigned a corresponding “local pointwise” topological entropy $h(P)$ [12] representing the connection map between distinct points of periodic orbit [13], and at the same time representing the infimum of the entropies of all corresponding maps which display orbits of the same configuration as P . On the other hand, for the piecewise monotone interval, the entropy of each continuous map f corresponds to the supremum of the number $h(P)$, relating in that way to sum of all periodic orbits P of f [14].

In this study we introduce certain situations where the non-local correlations arise from the non-Abelian structure of topological algebraic set systems, addressing at the same time the divergence in dimension and complexity measures of classical and quantum hypergraph representation. In particular, connection between the periodic orbits and the topological entropy for continuous maps on specific fractal hypergraph sets, realized by the action of the free group on two generators F_2 [15,16], is analyzed in Section 2. Geometric measure of exponential divergence between nearby orbits of F_2 free group is represented via fractional entropic descriptor [17] where action of two generators on a free group corresponds to iteration on two Lyapunov scales. In Section 3, we further exploit generalization of the Shannon measure entropy in order to assess relation between the Lyapunov exponents and the Rényi entropy [18], addressing at the same time some of the unique properties of the presented fractal set. The Rényi entropy as a measure of complexity of dynamical system is analyzed in a sense of correlation of probability distribution function to its local expansion rate, defined by the Lyapunov exponents. Dynamical features of trajectory based metric are further compared to topological entropy equivalent in order to fully quantify the parameters of complexity and characterize combinatorics of dynamical self-organization for the case of fractal system of free group on finite generators. Finally, in Sections 4 and 5, using the underlying fractal structure of the free group on two generators we construct the quantum hypergraph state [1] and demonstrate essential relation between its non-local correlations and the system complexity. Using the stabilizer formalism, the hypergraph states are defined as a class of multiqubit quantum states which generalizes graph states [19]. Obtained results allow efficient mapping of the quantum hypergraph states into corresponding logic circuits. Namely, in scope of the stabilizer formalism [20], in

terms of a group theory, it is possible to efficiently implement topological logic gates [21], using operations, such as controlled-NOT, Phase, Hadamard gates. In particular, a stabilizer code on n qubits (a quantum error-correcting code [22]) uses Pauli operators as stabilizer generators. For instance, in order to stabilize a subspace of 2^s dimension which belongs to the $2n$ dimensional Hilbert space, in total s stabilizer generators are required. In that case, $k = n - s$ logical qubits will be encoded into 2^{n-s} dimensional logical subspace. The space stabilized by the generators does not necessarily have to form an Abelian subgroup of the Pauli group over qubits [23]. In particular case of generators, where non-Abelian group is of size $s + 2e = n - k$ (k is notation of the number of qubits, codeword length is denoted as n ; e denotes ebits and s denotes the code ancilla qubit) the non-Abelian subgroup can be decomposed into two subgroups: the commuting isotropic group and the entanglement subgroup with anticommuting pairs [24].

2. Non-Abelian Statistics over Hypergraph Fractal Set

As we introduce some of the basic notations related to the hypergraph based fractal set, without loss of generality we assume topology embedded in \mathbb{R}^3 , where a hypergraph [25] represents a compact connected Hausdorff space represented by a subset $G(V, E)$ of vertex elements $V = \{v_1, v_2, \dots, v_n\}$ and edges $E \subset V$, $E \neq \emptyset$, where $E = \{e_1, e_2, \dots, e_m\}$. Considering a point $x \in V$, the number of edges which contain x will be denoted as the valence, v , of x . In case of a tree configuration where $v \geq 2$, we have the set of branching points $B(G)$ which coincide with the spatial hypergraph coordinates.

We shall first introduce some basic definitions related to group-theoretic construction of the specific graph sets. Thus, important property relating to the topological entropy of the special class of hypergraphs, introduced as a generalization of graphs in which infinite number of infinite clusters appears, is presented. Such geometric structure is the Cayley fractal set $\Gamma = \Gamma_{G,H}$ obtained by action of the free group G on finite generators, $H \subseteq G$.

Let G be a finitely generated group with the finite generating set $H = \{e_1, \dots, e_n\}$, where $H \subseteq G$. Then, the vertices $((x, u) \in \Gamma_p) \in G$ of corresponding sub-graph Γ_p , the set of edges $G \times \{e_1, \dots, e_n\}$ and corresponding mapping functions o and t where $o(x, e) = v$ and $t(x, e) = x \cdot e$, are all denoted by $\Gamma_{G,H}$. Action of the generator elements $e_i \in H$ over G connects vertices by inserting weighted edges $h = g \cdot e_i$ [26].

Consequently, a rank two free group, e.g., a free group G on two generators $H = (a^{\pm 1}, b^{\pm 1})$, $H \subseteq G$, forms a 4-valent tree. Detaching any edge from such 4-valent structure parts the underlying graph Γ_p into disjoint connected sets, resulting in infinitely many self-similar clusters for any p , where parameter $p \in [0, 1]$ determines connectivity of each edge of Γ_p , i.e., to be open with probability p and closed with probability $1 - p$. Uniqueness of the infinite cluster only appears when $p = 1$ and this property is connected to the notion of non-amenability [27], which directly relates to the non-Abelian statistics [28]. Hence, amenability in terms of a group structure directly influences power to determine a specific probability measure on G that is left invariant on subsets of G . Latter gives a clear relation between non-amenability and uniqueness of the fractal set, seen as an infinite cluster, which can be quantitatively assessed from the line of topological entropy and geometric concept of Lyapunov.

Example: Let $\Gamma = \Gamma_{G,H}$ be a Cayley tree where $\Gamma_p \subset \Gamma$ is its underlying graph structure. As a boundary of Γ_p we determine the edge-set $|\partial\Gamma_p|$ which represents the number of edges leaving Γ where $\partial\Gamma_p = \{(e \in x, u) | x \in \Gamma_p, u \in \Gamma \setminus \Gamma_p\}$.

Definition 1: Amenability [29], in the scope of the set theory, represents a property of possessing an invariant mean for the equivalence classes on almost all scale dimensions with respect to a given measure.

Definition 2: By the Følner condition [30] a set system defined on Cayley tree $\Gamma_{G,H}$ is amenable if and only if:

$$h(G) = \inf_{H \subset \Gamma} \frac{|\partial\Gamma_p|}{|\Gamma_p|} = 0, \tag{1}$$

where $|\cdot|$ denotes the related set cardinality, $\Gamma_p \subset \Gamma$ are finite subsets on a finitely generated group G , and $\partial\Gamma_p$ is the edge-set representing the boundary of Γ_p in reference to a given set of generators acting on G .

Definition 3—Dirichlet’s principle [31]: For each eventual partition of a set X (consisting of n elements, associated to positive integers) there exists a subset $x \subseteq X$, $|x|=N$, where N is a positive integer which corresponds to n , such that the mapping $\Pi : X \rightarrow X$ restricted to the subset x is either a constant or a 1-1 correspondence.

The above definition addresses existence and important relation of topological conjugacy between the set-system partitions and the elements of underlying subset.

Example: Let $T(X, E)$ be a tree with vertex set $X(T)$ and edge set $E(T)$. Partitioning of $T(X, E)$ into $P = \{X_1(T), X_2(T), \dots, X_n(T)\}$ corresponds to n sub-trees defined on a subset $x \subseteq X$, $|x|=N$, where the number of spanning trees $X_i(T)$ on N elements $x_i \subseteq X_i$ is given by a constant N^{N-2} [32].

Proposition 1: In case of a free group on two generators obtained tree structure $\Gamma_{G,H}$ is non-amenable, i.e., $h(G) \neq 0$ [33], because every step of selection over the tree (vertex) set $V(T) = \{X_i\}, i = 1, \dots, n$ (in order to form a connected sub-graph Γ_p) always generates the same number of boundary edges as vertices (due to the self-similar fractal property), i.e., any additional vertex element being appended to a connected sub-graph will create ≥ 2 additional edges $|\partial\Gamma_p|$.

Proof: (Erdos-Rado Canonization Lemma [31]) For each $X_i \in \mathcal{N}$ which is the generator of the i th tree on a sub-graph Γ_p , let the corresponding sub-graph vertex set $x(\Gamma_p) = \{x_i \subseteq X_i | x_i \in (x_1, \dots, x_N)\}$ be labeled with $x_1 = \{1\}, x_2 = \{1, 2\}, \dots, x_N = \{1, 2, \dots, N\}$. The order of Γ_p is $|x|=N$; it is associated to the number of x_i elements, which conform with the length- N Cayley permutation (C-permutation) p [34] of ordered set $x = \{x_1, \dots, x_N\}$, where $x_1 < x_2 < \dots < x_N$. Then, for every $p_x : |x|=N$ it follows that:

$$p_x = \frac{T_x}{\sum_{|x|=N} T_x}, \tag{2}$$

converges to a positive limit, where p_x is the weight probability distribution of each edge-set: $\partial\Gamma_p = \{(e_i \in x_i, u_i) | x_i \in \Gamma_p, u_i \in \Gamma \setminus \Gamma_p\}$, where two successive elements from x appear in C-permutation, defined for a stochastic sequence: $\{p^n\}_{n=1}^\infty$, $p^n = (p_1^n, p_2^n, \dots, p_N^n)$, and $T_x = e^{\lambda/\tau_x}$ is the probability distribution of the set $x = \{x_i\}$ on specified hypergraph partition, where $\sum_{|x|=N} T_x$ represents a cluster sum of distinguishable hypergraph partitions $\Pi: \{\sum_{|x|=N} T_x\}$ [35]. T_x is assigned on time interval: $0 < \tau_x \leq \infty$ which defines exponential growth where $\tau^* < \tau_x$ is the time of each vertex generation. T_x must be non-negative for each possible sequence of x . Thus, the random variable must have a positive value within the sequence of all $|x| = N$ possible values which are normalized to probability one, i.e., T_x must sum to one (directly addressing that $h(G) \neq 0$).

Consequently, the sequence $p^{n,max} := \max\{p_x : |x| = N\}$ converges to a positive limit, straightforwardly based on the stronger conjecture:

$p^{n,max}$ converges to a positive limit with probability one, where:

$$p^{n,max} = \frac{\max\{T_x : |x| = N\}}{\sum_{|x|=N} T_x}, \tag{3}$$

is the weight probability distribution of each edge-set: $\partial\Gamma_p = \{(e_i \in x_i, u_i) | x_i \in \Gamma_p, u_i \in \Gamma \setminus \Gamma_p\}$, where two successive elements from x appear in N -th level C-permutation for which: if $|x| = N$ (where $|x|$ is associated to the number of x_i elements), then the probability $T_x = e^{\lambda/\tau_x}$ is in total \sum_N -measurable (the sum is implicitly over all possible x_i) and as a result the denominator converges to a positive limit with probability one, $\sum_{|x|=N} T_x = 1$. Likewise, the numerator $T_x : |x| = N$ must converge to a positive limit with probability one for the maximal time interval $\tau_x \rightarrow \infty$, where $T_x = e^{\lambda/\tau_x} = 1$. \square

We focus next on a connection between uniqueness of the $\Gamma_{G,H}$ free group fractal set and the topological entropy, introducing a key role of non-amenability notion which is a prime characteristic of non-Abelian group topology [36]. Assuming a continuous map f of a compact metric space \mathcal{X} , the topological entropy represents the supremum of the metric entropy, where supremum is taken over all f -invariant Borel probability measures μ [37,38] on a topological space.

This property of the topological entropy can be easily extended for an arbitrary function $a(\cdot, t)$ which represents a scale if it is expanding for all intervals t and $\lim_{t \rightarrow \infty} a(s, t) = \infty$ for all scale parameters s . Hence, the topological entropy for a given growth scale a is:

$$h_{top}^a(f) = \sup_{\mu} h_{\mu}^a(f), \tag{4}$$

where supremum is taken over all localization parameters represented by the probability measures μ [39]. Now, for N -level Cayley tree $\Gamma_{G,H}$, where $\Gamma_p \subset \Gamma$ is its underlying graph structure, we define an f -invariant Borel probability measure μ on the finite generating set $x_i : |x| = N$, $x_i \in \Gamma_p$ (where $x = \{x_i \subseteq X_i | x_i \in (x_1, \dots, x_N)\}$ and $|x|$ represents the number of x_i elements, which give raise to the length- N Cayley permutation), as:

$$\mu(\{x\}) := \Delta_x, \tag{5}$$

where $\Delta_0 > 0$ and $\Delta_x = \sum_{k=1}^N \Delta_{x_k}$ are the probability distributions which denote the occurrences of the set system finite spatial partitions.

Example: Let mapping $f : M \rightarrow M$ define the metric space where $\lambda_1 > \lambda_2 > \dots > \lambda_k$ are the Lyapunov exponents of (f, μ) and E_k are the linear subspaces corresponding to exponents λ_k so that the dimension D of E_k corresponds to multiplicity of λ_k . For a hypergraph based fractal set of rank two free group, periodic orbits are defined on a hyperbolic $\varepsilon = 0$ space (due to a tree structure). In particular, geometric measure of the exponential divergence between nearby orbits of given sets is represented by the Lyapunov exponents, which can be successfully embedded into probabilistic framework in the following way.

Action of two generators on a free group corresponds to iteration of two scales which form the fractal set [40,41]. Correspondingly, two scales produce a spectrum of Lyapunov exponents: in this case for the symmetric map, $\lambda(p_x) = p_x \ln a + (1 - p_x) \ln b$, with $p_x = \frac{m}{n}$; $m, n \in \mathbb{Z}$ varying from 0 to 1 in the range: $0 \leq p_x \leq 1$ ($\lambda(p_x = 0) \leq \lambda_{\max}(p_x) \leq \lambda(p_x = 1)$), where $\lambda(p_x = 0) = \ln b$; and $\lambda(p_x = 1) = \ln a$.

In presence of the infinite sequences of intervals, which are defined by two scales $l_{m,n} = a^{-m} b^{-n+m}$ where $x = \frac{m}{n}$ is fixed, and n takes $\{n\}_1^\infty$ values, we have $N_n(\lambda_{\max}) = 2^n$ defined intervals and correspondingly $N_m(\lambda_{\min}) = n! / m!(n - m)!$ such defined intervals that determine the fractal dimension $D(\lambda_k)$ [17,40] as:

$$N_m = l_k^{-D(\lambda_k)}. \tag{6}$$

Here $l_k = e^{-k\lambda_k}$ relates directly to Lyapunov coefficients λ_k .

For the rank two generators $F_2 \{a^{\pm 1}, b^{\pm 1}\}$ action on a free group G , the corresponding Lyapunov coefficients are $\lambda_k = \frac{m}{n} \ln a^{\pm 1} + \frac{n - m}{n} \ln b^{\pm 1}$. Then, the entropic descriptor [17], *i.e.*, a measure of complexity associated to the fractal dimension $D(\lambda_k)$:

$$H(\lambda_k) = \log N_m \tag{7}$$

quantifies the number of unique sequences produced by the ratio $x = \frac{m}{n}$. Thus $x = (x_i)$ (according to $i = 1, \dots, N$ iterations) determines a periodic point [14] for the shift σ , where for each $n \geq 1$, there are 2^n points per period n for σ ; this specific property and its relation to topological entropy we discuss in the next section.

Now, the entropy H_n of the probability measure μ , which is defined on the finite (or countable) generating set $H\{x_i : |x| = N\}$, $H \subseteq G$ (where $\{\Gamma_p \subset \Gamma \mid x_i(\Gamma_p) \in (x_1, \dots, x_N)\}$ and $|x|$ is the number of x_i elements, *i.e.*, the order of Γ_p) for a Cayley fractal tree $\Gamma = \Gamma_{G,H}$ obtained after k permutations over elements of N -th sequence (x_N) , is given by:

$$\begin{aligned}
 H_n &= - \sum_{x_N=1}^k \Delta_{x_N} \log \Delta_{x_N} \\
 &= - \sum_{x_N=1}^k e^{\lambda_k / \tau_x} \log e^{\lambda_k / \tau_x} \\
 &= - \lambda_k / \tau_x \sum_{x_N=1}^k (e^{\lambda_k / \tau_x}),
 \end{aligned}
 \tag{8}$$

where $\sum_{x_N=1}^k \Delta_{x_N}$ are the probabilities for occurrence of the partitions: $\Pi_k = \left\{ \sum_{|x|=N} T_x \right\}$ that are produced by the generating set of $H \subseteq G$, where $T_x = e^{\lambda_k / \tau_x}$ is the probability operator acting over the specified generating set and λ_k is the Lyapunov coefficient.

2.1. Topological Entropy and Periodic Orbit Growth for Hypergraph Fractal Set

Without loss of generality, topological entropy is defined as a measure of maximal complexity of dynamical system.

Definition 4: Let \mathcal{A} be the finite set of n elements (alphabet), describing the discrete topology [42]. The dynamical system consisting of the sets of all bi-infinite symbol sequences $\mathcal{A}^{\mathbb{Z}} = \{x = (x_i)_{i \in \mathbb{Z}} : x_i \in \mathcal{A} \text{ for all } i \in \mathbb{Z}; i = 1, \dots, n\}$ represents the full \mathcal{A} -shift over a finite alphabet \mathcal{A} , where the shift map $\sigma : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$ shifts all sequences $x_i \in \mathcal{A}$ to the left, $(\sigma x)_i = x_{i+1}$.

Theorem 1: (Hasselblatt and Katok [43]) Topological entropy and periodic orbit growth coincide for shifts.

Proof: Let $\sigma : \sum_k \rightarrow \sum_k$ be a bilateral k -shift, acting on sequence $\{1, \dots, k\}$, and $\alpha = \{[1], \dots, [k]\}$ be a topological generator which transfers the spatial partitions of \sum_k into closed cylinders of length 1. Correspondingly, $\prod_{i=0}^{n-1} \sigma^{-i} \alpha$ denotes transformation of spatial partition of \sum_k into k^n cylinders of length n . As a result we have:

$$\begin{aligned}
 h_{top}(\sigma) &= h_{top}(\sigma, \alpha) \\
 &= \lim_{n \rightarrow \infty} \frac{1}{n} H_{top} \left(\prod_{i=0}^{n-1} \sigma^{-i} \alpha \right) \\
 &= \lim_{n \rightarrow \infty} \frac{1}{n} \log k^n,
 \end{aligned}
 \tag{9}$$

where for $n \rightarrow 1$ straightforwardly topological entropy for bilateral k -shift coincides with the logarithm of k sequences:

$$h_{top}(\sigma) = \log k.
 \tag{10}$$

Extension: $h_{top}(\sigma_k) = p(\sigma_k) = \log k$ stems from the restriction of the bilateral k -shift over invariant subset of k sequences, with a property of symbolic system [44]. □

The above can be easily explained using the following coin experiment.

Example: Let $\sigma, \sigma: M_0^\infty\{H, T\} \rightarrow M_0^\infty\{H, T\}$ be the twofold-switch operator acting over all spatial elements of $M_0^\infty\{H, T\}$ which denote the symbolic system (subshift) [44], where this operator determines the outcomes of an infinite series of trials (measurements). Then, considering that the sum of all possible outcomes in n trials is given by $N = \sum_n 2^n$, the topological entropy is defined as $h_{top}(\sigma) = \log 2$.

Thus, specifying the outcomes as $P(H) = p$ and $P(T) = 1 - p$ in fact determines the metric entropy $h_\mu(\sigma) = -p \log p - (1 - p) \log(1 - p)$. In case when the output is biased ($\{H, T\}$ represents the full shift), the number of typical outcomes in n trials is $\sim e^{nh}$ for $h < \log 2$, representing in that way the inequality relation between the topological $h_{top}^\sigma(f)$ and the metric entropy $h_\mu^\sigma(f)$:

$$h_{top}^\sigma(f) \leq \sup_\mu h_\mu^\sigma(f). \tag{11}$$

Lemma 1: Extending relation (10) to a hypergraph subset $G(V, H)$ for a finite type subshift [44]: $\sigma: \sum_{M_{i,j}} \rightarrow \sum_{m_{i,j}}$ where M is irreducible $n \times n$ transition matrix with entries in $\{0, 1\}$, under condition that the largest eigenvalue is always real by the Perron-Frobenius theorem [45], results that the topological entropy $h_{top}(\sigma)$ is equal to the largest positive eigenvalue of matrix M .

Proof: Let $\alpha = \{[1], \dots, [n]\}$ be a topological generator which induces partitions $\bigvee_{i=0}^{n-1} \sigma^{-i} \alpha$ of space $\sum_{M_{i,j}}$ into a closed cylinders of length n , where σ is a subshift of finite type, and resulting topological entropy coincides with the number of cylinders of length n in $\sum_{M_{i,j}}$

$$H_{top} \left(\bigvee_{i=0}^{n-1} \sigma^{-i} \alpha \right) = \log \text{card} \left\{ \text{no. of cylinders of length } n \text{ in } \sum_{M_{i,j}} \right\}, \tag{12}$$

where cylinder set of length n is defined on space $[i_0, \dots, i_{n-1}] \cap \sum_{M_{i,j}} \neq \emptyset$ if and only if $M_{i_0, i_1} M_{i_1, i_2} \dots M_{i_{n-2}, i_{n-1}} = 1$ and the number of length- n cylinders that overlap space $\sum_{M_{i,j}}$ is $\|M^n\|$.

As a result, the connection between the topological entropy and the transition matrix [15] is given by:

$$\begin{aligned} h_{top}(\sigma) &= h_{top}(\sigma, \alpha) \\ &= \frac{1}{n} H_{top} \left(\bigvee_{i=0}^{n-1} \sigma^{-i} \alpha \right) \\ &= \frac{1}{n} H_{top} \|M^n\| \\ &= \log \lambda_{i_{max}}, \quad \square \end{aligned} \tag{13}$$

where the topological entropy equals to the logarithm of the largest eigenvalue of the transition matrix (spectral radius) [46].

2.1. Relations between Lyapunov Spectrum, System Dimension and Measure Entropy for Hypergraph Fractal Set

In case of the systems with infinite degrees of freedom (e.g., the presented fractal system of the free group on two generators, where applies the notion of non-amenability given by Equation (1)) assuming d -dimensional system of linear size L_d , it is possible to establish relation between Lyapunov spectrum, dimension of the system and measure entropy. The first step towards such relation is to resolve the case of 'thermodynamic' limit, $L \rightarrow \infty$, for the Lyapunov spectrum, which can be assessed by analyzing whether ratio $\lambda_i : x = i/L_d, x \in [0, 1]$, converges to a density function $\lambda_i = \Lambda(x)$ as $L \rightarrow \infty$.

Starting from the Pesin formula [47], where θ is the Heaviside function:

$$h = \sum_i \lambda_i \theta(\lambda_i), \tag{14}$$

which infers that the Kolmogorov-Sinai entropy [48,49] is in this case proportional to the maximum Lyapunov exponent $h = \sum_{i=1}^p \lambda_i$. Thus, there is only one positive Lyapunov exponent, as the system size approaches thermodynamic limit $L \rightarrow \infty$.

Local entropy h_i follows from Equation (14); it is defined for each degree of freedom for the system of linear size L_d , as:

$$h_i = \lim_{L \rightarrow \infty} \frac{h}{L_d} = \int_0^1 \Lambda(x) \theta(\Lambda) dx, \lambda_i = \Lambda(x), \tag{15}$$

where after setting the existence of bound: $\lambda_i = \Lambda(x)$ as $L \rightarrow \infty$, total entropy h is directly related to the dimension of the system. Indeed, using the Kaplan and Yorke formula [50], the fractal dimension D can be estimated from the Lyapunov spectrum as:

$$D = p + \frac{\sum_{i=1}^p \lambda_i}{|\lambda_{p+1}|}. \tag{16}$$

Also, under condition: $\sum^p \lambda_i > 0$, the dimension of the attractor, D_λ , is proportional to L_d leading to the existence of inherent dimension density $\delta_\lambda = \frac{D_\lambda}{L_d}$ per system degree of freedom.

Now we are ready to elucidate the essential connection between the Lyapunov exponents λ_i and d -regular hypergraph fractal set. We use the set-theoretic approach, starting from the structure of the Cayley tree. The Cayley tree on a free group G generated by the finite set $H \subseteq G$ is represented by the graph with vertex set $V \in G$ and the edge set $\{(a, b) | a, b \in H\}$. Let G be a free group on two generators where $L_2(G)$ is the function space on G , then $L_2(G)$ can be decomposed into subspaces: $L_2(G) = \bigoplus_{i=1}^r E_i$ meaning that it has the value 1 on the i -th vertex of a set G and 0 otherwise.

Proposition 2: Assume that the Cayley set satisfies relation: $H = H^{-1}$. This condition as a result infers existence of orthogonal set of eigenvectors and eigenvalues belonging to the adjacency matrix A . Under constraint that the eigenvalues of A , i.e., $\lambda_1, \dots, \lambda_{d_i}$ are bounded by corresponding vector subspaces E_i , the norm of the F -uniform, d -regular (each vertex has degree d) Cayley hypergraph on n vertices is defined as $\left(\frac{n}{d_i}\right) (F-2)/2 \max_i(\lambda_i)$ where $\max_i(\lambda_i) \in \lambda_{i,i} = \frac{d}{n} M_{i,i}$, and $M_{i,i}$ is a diagonal matrix.

Proof: We first demonstrate that $H = H^{-1}$ property directly implies that M as a symmetric and a real matrix can be diagonalized by a set of eigenvectors in \mathbf{R}^d . Namely, if H is a Hermitian matrix, then $H = UMU^H$, where U is unitary matrix and M is a diagonal matrix (and as such also a symmetric matrix) with real entries λ_i . Therefore: $H^{-1} = (UMU^H)^{-1} = (U^H)^{-1} M^{-1} U^{-1} = UM^{-1}U^H$, since $U^{-1} = U^H$. Here M^{-1} is a diagonal matrix with corresponding elements $1/\lambda_i$. Likewise, H^{-1} is also a Hermitian matrix where $(H^{-1})^H = (UM^{-1}U^H)^H = U(M^{-1})^H U^H = UM^{-1}U^H = H^{-1}$. Now, let A and F be the adjacency matrix and generating function of the Cayley hypergraph, respectively. A and F are bounded on the vector subspace E_i corresponding to the d dimensional state ρ (where the coefficients of $\{\rho_{i,j}\}$ are given with respect to elements of the basis E_i) as:

$$\sum_{x,y \in H} \rho_{i,j}(x) \rho_{k,l}(y) = \sum_{g \in G} \rho_{i,j}(g) \rho_{k,l}(g^{-1}h), \tag{17}$$

leading to:

$$\sum_{g \in G} \sum_{m=1}^d \rho_{i,j}(g) \overline{\rho_{l,m}(g)} \rho_{m,k}(h) = \delta_{i,k} \left(\frac{n}{d}\right) \rho_{j,l}(h), \tag{18}$$

where $\delta_{i,k}$ is the Kronecker delta function, ensuing that generating function satisfies:

$$F\left(\sum_{i,j} \alpha_{i,j} \rho_{i,j}, \sum_{k,l} \beta_{k,l} \rho_{k,l}\right) = \sum_{i,j,k,l} \alpha_{i,j} \beta_{k,l} \delta_{i,k} M_{j,l}, \tag{19}$$

where $M_{j,l}$ are elements of the diagonal matrix:

$$M = \left(\frac{n}{d}\right) \sum_{h \in H} \rho(h). \tag{20}$$

M acts as a diagonal matrix by extending these eigenvectors to the complex basis of C^d . Consequently, as the normalization for each $\rho_{i,j}$ element is given by $\sqrt{n/d}$ it follows that for each i, j the coefficients $\rho_{i,j}$ represent eigenvectors of the adjacency matrix A with eigenvalue 0 if $i \neq j$, and eigenvalue $\lambda_{i,i} = \frac{d}{n} M_{i,i}$ if $i = j$.

Now we can easily generalize the concept of the Cayley $\Gamma = \Gamma_{G,H}$ graph to the corresponding hypergraph set.

Proposition 3: The eigenvalues of the d -regular Cayley hypergraph can be determined from the $\Gamma = \Gamma_{G,H}$ subgraph and from the $L_2(G)$ decomposition.

Proof: Let H be the symmetric set of generators with the vertex set assigned on a finite group G , then the d -regular Cayley hypergraph on G and H is composed of edges $E = \{(e_1, \dots, e_d) : e_1 \dots e_d \in H\}$. Because of its tree character, d -regular Cayley hypergraph has intrinsic Markov property and it can be analyzed in terms of a piecewise linear Markov map f [51] according to the spectral radius of matrix M (whose elements are $\{0, 1\}$). Direct consequence is that the topological entropy corresponds to the maximal eigenvalue of the matrix M as it is already shown by Equation (13). \square

Moreover, taking into account the bound $\lambda_i = \Lambda(x)$ as $L \rightarrow \infty$ where $\lambda_{i\max} = \max_i \sqrt{\frac{n}{d}} \lambda_{i,i}$, it follows that Lyapunov spectrum equals to the maximal Lyapunov exponent $\lim_{L \rightarrow \infty} \Lambda(x) = \max_i \sqrt{\frac{n}{d}} \lambda_{i,i} = \sum_{i=1}^p \lambda_i$. As a result the Kolmogorov-Sinai entropy in case of d -regular Cayley hypergraph is proportional to:

$$h = \sum_{i=1}^p \lambda_i = \max_i \sqrt{\frac{n}{d}} \lambda_{i,i}. \tag{21}$$

3. Rényi Topological Entropy and Lyapunov Exponents for Non-Abelian Fractal Set

Assume that the fractal set is partitioned into distinguishable clusters of size r . In order to localize a point with a precision r one must determine the partition which contains the point and quantify average data on particular partition. For this one can use the Shannon information formula, and also the topological Rényi entropy of order q , which is directly related to the generalized dimension [52]:

$$D_q = \lim_{r \rightarrow 0} \frac{\sum_i P_i \log P_i}{\log r} = \frac{1}{q-1} \lim_{r \rightarrow 0} \frac{\log \sum_i P_i^q}{\log r}, \tag{22}$$

where P_i is the probability measure of the i th partition.

Now, let Π be a partition assigned by a free group G on rank two generators $H \subseteq G$ where $\{\pi_i^{(n)}\}$ are the elements of partitions Π^n obtained under measure preserving transformation [53]:

$$\Pi_n = \bigvee_{i=0}^{n-1} f^{-i}(\Pi). \tag{23}$$

The Rényi entropy of order q is the supremum over all distinguishable partitions Π^n :

$$H(q) = \sup_{\Pi} \left\{ \lim_{n \rightarrow \infty} \frac{1}{1-q} \frac{1}{n} \ln \sum_i \mu(\Pi_i^{(n)})^q \right\}. \tag{24}$$

Next step establishes relation between the topological Rényi entropy of order q and the generalized Lyapunov exponents Λ . By assigning the probability distribution function $p_x(i_0 \dots i_{n-1})$ to the

occurrence of distinguishable partitions Π^n that are specified in time steps: $i_0(t=0), i_1(t=1), \dots, i_{n-1}(t=n-1)$ as $\sum_{i_0 \dots i_{n-1}} (p_x(i_0 \dots i_{n-1}))^q$ for which:

$$H(q) = \lim_{\varepsilon \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{1-q} \frac{1}{n} \ln \sum_{i_0 \dots i_{n-1}} (p_x(i_0 \dots i_{n-1}))^q, \tag{25}$$

and using the generalized measure $\Lambda(i_k)$ which represents the average expansion rate of information given by the building block i_k of dimension ε which constituents each partition Π_i ; the initial building block i_0 is defined by the corresponding Lyapunov spectrum $\Lambda(i_0)$. After the initialization is performed, the Lyapunov spectrum is given by: $\Lambda(i_0)\Lambda(i_1), \dots, \Lambda(i_{n-1})$ from which we obtain the probability rate for the expansion of the spatial partitions building blocks:

$$\sum_{i_0 \dots i_{n-1}} p_x(i_0 \dots i_{n-1}) = \frac{\mu(i_0)}{\Lambda(i_0) \dots \Lambda(i_{n-1})}. \tag{26}$$

Substitution of Equation (26) into Equation (25) gives:

$$H(q) = \lim_{\varepsilon \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{1-q} \frac{1}{n} \ln \left[\alpha_n \left\langle \left(\frac{\mu(i_0)}{\Lambda(i_0) \dots \Lambda(i_{n-1})} \right) \right\rangle^q \right], \tag{27}$$

where α_n is obtained from the normalization: $\sum_{i_0 \dots i_{n-1}} p_x(i_0 \dots i_{n-1}) = 1$ and represents the number of various distinguishable clusters (blocks) of length n . Now, from Equation (27) follows the measure:

$$H(q) = \lim_{\varepsilon \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{1-q} \frac{1}{n} \ln \left[\left\langle \frac{\mu(i_0)}{\Lambda(i_0) \dots \Lambda(i_{n-1})} \right\rangle^{-1} \left\langle \left(\frac{\mu(i_0)}{\Lambda(i_0) \dots \Lambda(i_{n-1})} \right) \right\rangle^q \right], \tag{28}$$

which establishes the correlation between the topological Rényi entropy of order q and the generalized Lyapunov exponents $\Lambda(i_k)$, for the limit $n \rightarrow \infty$ this correlation reads:

$$H(q) = \lim_{\varepsilon \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{1-q} \frac{1}{n} \ln \left\langle \exp \left((1-q) \sum_{k=0}^{n-2} \ln \Lambda(i_k) \right) \right\rangle. \tag{29}$$

Example: Let $G(V, H)$ be a hypergraph defined on a free group G on finite generators H , where vertex set: $V_i = \{x_1, x_2, \dots, x_n\}_1^n \in G, H \subseteq G$, and topological order coefficients: $q_i = (q_1, \dots, q_n), \sum_{x \in H_i} q_i \geq 1$, are assigned on all local partitions Π_i , such that:

$$\begin{aligned} q_1 &\rightarrow \Pi_1 \left(\{p_{x_1}, p_{x_2}, \dots, p_{x_n}\}_1^n \right), q_2 \rightarrow \Pi_1 \left(\{p_{x_1}, p_{x_2}, \dots, p_{x_n}\}_1^n \right) \cdot \Pi_2 \left(\{p_{x_1}, p_{x_2}, \dots, p_{x_n}\}_1^n \right), \dots, \\ q_n &\rightarrow \Pi_1 \left(\{p_{x_1}, p_{x_2}, \dots, p_{x_n}\}_1^n \right) \cdot \Pi_2 \left(\{p_{x_1}, p_{x_2}, \dots, p_{x_n}\}_1^n \right) \cdot \dots \cdot \Pi_n \left(\{p_{x_1}, p_{x_2}, \dots, p_{x_n}\}_1^n \right), \end{aligned}$$

and defined on a metric space M_x with probability measure μ , where a nonnegative function $f_i : \prod_{x \in H_i} M_x \rightarrow \mathbf{R}$ is associated to each generator set H_i . Then, under constraint of integrability [38]:

$$\int \prod_i f_i \prod_{x \in V_i} d\mu_x \leq \prod_i \left(\int f_i^{1/q_i} \prod_{x \in H_i} d\mu_x \right)^{q_i}, \tag{30}$$

and according to Equation (24) where the Rényi entropy of order q is the supremum over all distinguishable hypergraph partitions, it follows the expression for the fractional Rényi entropy of order q for a hypergraph defined on a free group G with respect to the generator set $H_i \subseteq G$:

$$H(q) = \frac{1}{1-q} \log \int_x \prod_i \left(\int f_i^{1/q_i} \prod_{x \in H_i} d\mu_x \right)^{q_i}. \tag{31}$$

In particular case of two generators, the Rényi's entropy dependence from the probability measure μ_x , and the probability rate for the expansion of the spatial partitions (Equation (31)), for different values of order q , is presented in Figure 1 and Figure 2, respectively. Figure 3 shows the maximum allowed correlation between the topological Rényi entropy of order q and the generalized Lyapunov exponents $\Lambda(i_k)$ for the partitioned hypergraph fractal set (Equation (24) obtained by action of the rank -2 free group on finite generators [26].




Figure 1. Topological Rényi entropy as a function of the probability measure μ_x over hypergraph fractal set of free group on finite generators, according to equation (31), and order: $q = 0.4$ (dash-dotted), $q = 5$ (dashed) and $q = 1$ (full line).




Figure 2. Plot of the output entropies as functions of probability rate for the expansion of the spatial partitions (Equation (31)) for different input order parameters (designated by solid lines): $q = 0.2$ (blue), $q = 0.4$ (yellow), $q = 0.5$ (dark cyan), $q = 0.8$ (dark yellow).

4. Underlying Non-Locality of F_2 —Hypergraph State

In this section we present a more structural understanding of non-locality [54] of a hypergraph state which is obtained by action of the rank two free group, called F_2 . In particular, we show that intrinsic non-local character of the non-Abelian actions [55] of the free group on two generators is closely tied to notion of the non-amenability.

We consider the algebraic topological structure obtained via generating set HUH^{-1} , $H := \{a^{\pm 1}, b^{\pm 1}\}$, where $a; b$ denote basis of the free group F_2 on two generators $\{a^{\pm 1}, b^{\pm 1}\}$ which realize an infinite 4-regular tree represented as the undirected Cayley set. In this case, the infinite 4-regular structure gives a hypergraph based covering space for the wedge of two circles $H^1 \wedge H^1$, $(H_{\pm}^1 = a^{\pm 1} + b^{\pm 1} = 1)$ which has fundamental ($\delta = 0$) hyperbolic group $F_2 \cong \mathbb{Z} * \mathbb{Z}$ where $\pi_1(H^1 \wedge H^1, 0) \cong \pi_1(H^1, 0) * \pi_1(H^1, 0)$.

Definition 5: Group G is amenable if and only if it does not produce paradoxical decomposition [43].

From the constraint of amenability and Equation (1) it follows that a rank two free group $F_2 = a, a^{-1}, b, b^{-1} | a, b \in H$ of $SO(3)$ is non-amenable and as a result it is non-Abelian [27] (see fundamental theorem of finitely generated Abelian groups), and there exists a countable subset H on the sphere S such that the decomposition $S - H$ is F -paradoxical [56,57]; a straight repercussion is that $SO(3)$ is paradoxical too. Thus, the elements of the free group of two generators $F_2 \in SO(3)$ are

distance preserving as operators on \mathbb{R}^3 , where they represent a group of nontrivial, independent rotations of the sphere about axis that passes through the sphere center.




Figure 3. Maximum allowed correlation between the topological Rényi entropy of order q and the generalized Lyapunov exponents $\Lambda(i_k)$ obtained for the supremum output entropy conjecture over all distinguishable partitions (Equation (24)). The x - y axes are the average expansion rate of information given by the building block i_k of dimension ε on N intervals and input order parameters q , respectively.

Likewise, there are rotations a_R, b_R of the unit sphere in \mathbb{R}^3 that generate exactly the free group on two generators. To prove that let us define a group of rotations $(a_R(\varphi), b_R(\varphi))$ for $\varphi = \arccos(1/3)$ about orthogonal axes:

$$a_R(\varphi) = \begin{bmatrix} 1/3 & -2\sqrt{2}/3 & 0 \\ 2\sqrt{2}/3 & 1/3 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad b_R(\varphi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/3 & -2\sqrt{2}/3 \\ 0 & 2\sqrt{2}/3 & 1/3 \end{bmatrix}. \tag{32}$$

In this case elements of a group: $a_R; b_R; a_R^{-1}; b_R^{-1}$ map vector basis $(1, 0, 0)$ to a basis $(x, y\sqrt{2}, z)/3^n$, where x, y , and z as integers, $y \neq 0$, and n is the length of the information string.

Proposition 4: Let $f : a_R \rightarrow a, f : b_R \rightarrow b$, then for a generating set $H : \{a, b\}$ we can define a rank 2 free group $F_2 = a, a^{-1}, b, b^{-1} \mid a, b \in H$, which determines paradoxical decomposition of $S - H$, manifesting in

this way intrinsic non-local feature of the hypergraph fractal state based on F_2 free group. In particular this means that $F_2 \in SO(3)$ is partitioned into five parts where only four are needed to establish two exact copies of the original sphere S , *i.e.*, to perform the paradoxical decomposition.

Proof: The natural decomposition of $F_2 \in SO(3)$ is given by:

$$F_2 = \{e\} \cup \omega(a) \cup \omega(a^{-1}) \cup \omega(b) \cup \omega(b^{-1}), \tag{33}$$

where for all disjoint sets, $\omega(\cdot)$ denotes all possible information strings starting with (\cdot) .

The following relations establish correlation between information strings and the set of free group on two generators:

$$\begin{aligned} \omega(a^{-1}) &= a^{-1}(F_2 \setminus \omega(a)), \\ \omega(b^{-1}) &= b^{-1}(F_2 \setminus \omega(b)). \end{aligned} \tag{34}$$

But, another decomposition yields the following property:

$$F_2 = \omega(a) \cup \omega(a)^c = \omega(a) \cup a\omega(a^{-1}), \tag{35}$$

where $\omega(\cdot)^c$ is the complement of the corresponding set $\omega(\cdot)$, and where actual multiplying $\omega(a^{-1})$ by a acts like translation over $\omega(b)$ and $\omega(b^{-1})$. Here $h \in F_2 \setminus \omega(a)$, then $a^{-1}h \in \omega(a^{-1})$, and $h = a(a^{-1}h) \in a\omega(a^{-1})$.

Correspondingly, the third decomposition of F_2 is given by:

$$F_2 = \omega(b) \cup \omega(b)^c = \omega(b) \cup b\omega(b^{-1}). \tag{36}$$

Now, in order to demonstrate the notion of non-amenability which directly implicates non-locality of the hypergraph quantum states based on the free group F_2 , one can start by assigning the mean probability distribution $m(F_2) = 1$ that maps subsets of $F_2 = a, a^{-1}, b, b^{-1} | a, b \in H$ to the unit interval $[0, 1]$ as following:

$$\begin{aligned} m(F_2) = 1 &= m(\omega(a)) + m(a\omega(a^{-1})) \\ &= m(\omega(a)) + m(\omega(a^{-1})), \end{aligned} \tag{37}$$

likewise:

$$\begin{aligned} m(F_2) &= 1 \\ &= m(\omega(b)) + m(\omega(b^{-1})). \end{aligned} \tag{38}$$

Then, from Equation (33) we have:

$$m(F_2) = m(\{1\}) + m(\omega(a)) + m(\omega(a^{-1})) + m(\omega(b)) + m(\omega(b^{-1})). \tag{39}$$

Comparison of previous relation with Equations (37) and (38) directly gives:

$$\begin{aligned}
 & m(\{1\}) + m(\omega(a)) + m(\omega(a^{-1})) + m(\omega(b)) + m(\omega(b^{-1})) \\
 & \geq m(\omega(b)) + m(\omega(b^{-1})) + m(\omega(a^{-1})) + m(\omega(a)).
 \end{aligned}
 \tag{40}$$

From Equations (37), (38) and (40) we have:

$$\begin{aligned}
 & m(\omega(a)) + m(\omega(a^{-1})) = m(\omega(b)) + m(\omega(b^{-1})) \Rightarrow \\
 & m(\{1\}) + m(\omega(a)) + m(\omega(a^{-1})) + m(\omega(b)) + m(\omega(b^{-1})) \geq 2(m(\omega(b)) + m(\omega(b^{-1}))).
 \end{aligned}
 \tag{41}$$

Taking into account equation (40), latter also holds for:

$$\begin{aligned}
 & m(\{1\}) + m(\omega(a)) + m(\omega(a^{-1})) + m(\omega(b)) + m(\omega(b^{-1})) \\
 & \geq m(\omega(b)) + m(\omega(b^{-1})) + m(\omega(a)) - m(\omega(a^{-1})).
 \end{aligned}
 \tag{42}$$

Thus, from Equations (37), (38) and right side of Equation (42), directly follows

$$m(\omega(b)) + m(\omega(b^{-1})) + m(\omega(a)) - m(\omega(a^{-1})) \leq 2\sqrt{(\omega(a))^2}, \tag{43}$$

finally, by inserting into Equation (43) the relations which address the mean probability distribution of F_2 :

$$F_2 = 1 = \omega(a) + \omega(a^{-1}) \Leftrightarrow p + (p-1),$$

where:

$$\begin{aligned}
 \omega(a) &= 1 - \omega(a^{-1}), \\
 &= 1 + (1 - p),
 \end{aligned}$$

we obtain the characteristic inequality relation for the free group F_2 set corresponding to CHSH inequality [58]:

$$CHSH = m(F_2)_{CHSH} \leq 2\sqrt{(\omega(a))^2} \approx 2\sqrt{(1+(1-p))^{2N}}, \quad 0 \leq p \leq 1. \tag{44}$$

where N is the number of sequences of intervals corresponding to the fractal dimension (see Figure 4.) where uniqueness of the infinite cluster only appears when $p = 1$. Analogue inequality relation can be obtained for all corresponding decomposition elements of $F_2 \in SO(3)$. Former contraintuitive inequality [54,58] directly provides the resource for analysis of the non-local properties of the hypergraph quantum states [1], defined in this case on the 4-regular subset $a, a^{-1}, b, b^{-1} | a, b \in H$ of the free group, F_2 . Correlation over the elements of the mean probability distribution of F_2 corresponds to operations of translation and inversion between information strings starting with: $C(\omega(a), \omega(b)), C(\omega(a), \omega(b^{-1})), C(\omega(a^{-1}), \omega(b)), C(\omega(a^{-1}), \omega(b^{-1}))$. In that context, bound $p = 1$ on $m(F_2)_{CHSH}$ (Equation (44)) establishes demarcation line between the complexity of the classical (local realism) and the quantum counterpart [59] of system state.

5. Quantum Hypergraph State

First step towards obtaining the quantum hypergraph state [1, 19, 60] from a mathematical hypergraph $G(V, H)$ based on the free group F_2 , is to assign qubits to a vertex set elements, $V = \{x_1, x_2, \dots, x_n\}$, and next is to perform initialization of qubits into $|+\rangle$ state [60, 61]. Subsequent control and manipulation over qubit states is achieved through hyperedge controlled-Z operations [1, 60] between specified qubits, which position coincides with underlying geometry of the free group- F_2 fractal set. G is the free group on two generators which form a set $H = (a^{\pm 1}, b^{\pm 1})$, $H \subseteq G$, where every element $g \in G$ can be obtained by the action HUH^{-1} . As a result, a 4-valent hypergraph is generated by action $H \times G$. Qubit states are assigned as:

$$\{a, a^{-1}\} = \{|0\rangle|0\rangle, |1\rangle|0\rangle\}, \quad \{b, b^{-1}\} = \{|0\rangle|0\rangle, |1\rangle|0\rangle\}, \tag{45}$$

where measurement basis is specified in exact correspondence to a rank two generating set on a free group $F_2 = a, a^{-1}, b, b^{-1} | a, b \in H$, where four basis states are: $O = \{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$.




Figure 4. Maximum violation of $m(F_2)_{CHSH}$ inequalities obtained for $S_{\max} \approx 2.82$ and $p \geq 0$ versus number of interval sequences N (Equation (44)), corresponding to CHSH inequality [58,59], plotted for four order parameters (designated by solid squares): $q = 0.2$ (blue), $q = 0.4$ (dark yellow), $q = 0.5$ (dark cyan), $q = 0.8$ (cyan).

The main prerequisite for encoding is to perform the initialization of the system with probabilities

$$\mathcal{M}_0^e = \{p_1 = 1000, p_2 = 0100, p_3 = 0010, p_4 = 0001\}, \quad \sum_{i=1}^n \left(\sum_{k=1}^4 p_k \right)_i = 1.$$

In the first iteration, the probability distribution arising from the corresponding set of measurements \mathcal{M}_1 over the Cayley subset $\{\{a, b\}, \{a, b^{-1}\}, \{a^{-1}, b\}, \{a^{-1}, b^{-1}\}\}$ is given by:

$$\mathcal{M}_1 = \{\{a, b\}, \{a, b^{-1}\}, \{a^{-1}, b\}, \{a^{-1}, b^{-1}\}\} :$$

$$\{\{0000, 0100, 1000, 1100\}, \{0001, 0101, 1001, 1101\}, \{0010, 0110, 1010, 1110\}, \{0011, 0111, 1011, 1111\}\}^{i=\pm 1} ,$$

where action of the measurement is determined by the projective measurement operator $Pi_X = |X\rangle\langle X|$ over a given generating set (see Equation (46)), producing in that way the state ρ which possesses non-local correlations, representing at the same time the topological state. In particular, the symmetry of the underlying topology is efficiently captured and corresponds exactly to the symmetry of the joint probabilities for each measurement, see Figure 5. In this case the tensor algebra $T(V) = \bigoplus_{i \geq 0} V^{\otimes i}$ is represented on a k -module V of a vertex set elements, which is an associative k -algebra spanned by $x_1 x_2 \dots x_n := x_1 \otimes x_2 \otimes \dots \otimes x_n$, where $n \in \mathbb{N}$ and $x_1, x_2, \dots, x_n \in V$. Multiplication is utilized by a k -linear map $m(x_1 x_2 \dots x_n \otimes \omega_1 \omega_2 \dots \omega_m) := x_1 x_2 \dots x_n \omega_1 \omega_2 \dots \omega_m$ for all $n, m \in \mathbb{N}$ and $x_1, x_2, \dots, x_n, \omega_1, \omega_2, \dots, \omega_m$ in V .

$$\left[\begin{array}{c} \boxed{\begin{array}{c} |0\rangle|0\rangle \\ |1\rangle|0\rangle \end{array}}_a \\ \boxed{\begin{array}{c} |0\rangle|1\rangle \\ |1\rangle|1\rangle \end{array}}_{a^{-1}} \end{array} \right] \otimes \left[\boxed{\begin{array}{c} |0\rangle|0\rangle \quad |1\rangle|0\rangle \end{array}}_b \quad \boxed{\begin{array}{c} |0\rangle|1\rangle \quad |1\rangle|1\rangle \end{array}}_{b^{-1}} \right] = \left[\begin{array}{cc} \boxed{\begin{array}{c} |00\rangle|00\rangle \quad |01\rangle|00\rangle \\ |10\rangle|00\rangle \quad |11\rangle|00\rangle \end{array}}_b & \boxed{\begin{array}{c} |00\rangle|01\rangle \quad |01\rangle|01\rangle \\ |10\rangle|01\rangle \quad |11\rangle|01\rangle \end{array}}_a \\ \boxed{\begin{array}{c} |00\rangle|10\rangle \quad |01\rangle|10\rangle \\ |10\rangle|10\rangle \quad |11\rangle|10\rangle \end{array}}_{a^{-1}} & \boxed{\begin{array}{c} |00\rangle|11\rangle \quad |01\rangle|11\rangle \\ |10\rangle|11\rangle \quad |11\rangle|11\rangle \end{array}}_{b^{-1}} \end{array} \right] \quad (46)$$




Figure 5. (a) Geometrical-representation of a rank-2 free group $F_2 = a, a^{-1}, b, b^{-1} \mid a, b \in H$, where $H \subseteq G$ is a generating set. (b) Scheme of the hypergraph state quantum correlations [1,60,62] that reflect exact symmetry of the underlying $F_2 = a, a^{-1}, b, b^{-1} \mid a, b \in H$ topology, (Equation (46)).

Likewise, due to a self-similar property, in each subsequent action performed over d -dimensional Cayley fractal structure that corresponds to a rank- d generating set on a free group F_d , probability distributions arising from the corresponding set of measurements determined by the subset H are

$$\mathcal{M}_n = \left\{ \{a^n, b^n\}, \{a^n, b^{-n}\}, \{a^{-n}, b^n\}, \{a^{-n}, b^{-n}\} \right\} : \\ \left\{ \{0000, 0100, 1000, 1100\}, \{0001, 0101, 1001, 1101\}, \{0010, 0110, 1010, 1110\}, \{0011, 0111, 1011, 1111\} \right\}^{i=\pm n},$$

Bounds of non-local correlations [63] are closely assessed by determination of observables that influence the maximum of $I \sim \lambda_{i, \max}$ where $I = \text{Tr}(\mathcal{B}\rho)$ [62] with \mathcal{B} denoting the Bell operator, and by eigenvalues $\lambda_{i, \max}$ which determine violation of Bell inequality for the concrete joint probabilities.

$\{\widehat{a}_{A_k}\}, \{\widehat{b}_{B_k}\}, \{\widehat{a}_{A_k}^{-1}\}, \{\widehat{b}_{B_k}^{-1}\}$ are the orthonormal eigenvectors of the corresponding observables $\{A_1, \dots, A_n\}$ and $\{B_1, \dots, B_n\}$ defined on a vertex set $V = \{x_1, x_2, \dots, x_n\}$ of the free group $G(V, H)$ on two generators, F_2 . Consequently, for the d -regular hypergraph state on n vertices, the maximal value for I is bounded with the maximal eigenvalue $\lambda_{i, \max} = \max_i \sqrt{\frac{n}{d}} \lambda_{i, i}$ of the corresponding diagonal element of $d \times d$ matrix U_ρ :

$$U_\rho = \begin{pmatrix} \lambda_{0,0} & \cdots & \lambda_{0,d-1} \\ \vdots & \ddots & \vdots \\ \lambda_{d-1,0} & \cdots & \lambda_{d-1,d-1} \end{pmatrix}, \tag{47}$$

obtained with operator $U = \left[\prod_{i=0}^{d-2} \left(\prod_{j=i+1}^{d-1} e^{(iP_j \lambda_{j,i})} e^{(i\sigma_{i,j} \lambda_{i,j})} \right) \right] \left[\prod_{k=0}^{d-1} e^{(iP_k \lambda_{k,k})} \right]$, where $\sigma_{i,j} = -i|i\rangle\langle j| + i|j\rangle\langle i|$, and off-diagonal elements $\lambda_{i,j}, \lambda_{j,i}$ denote rotation and a phase shift, respectively. For the case when $A_k = \sigma_i, A_k^{-1} = \sigma_j, B_k = \frac{1}{\sqrt{2}}(\sigma_i + \sigma_j), B_k^{-1} = \frac{1}{\sqrt{2}}(\sigma_i - \sigma_j); k = 1, \dots, n$, Bell operator, \mathcal{B} , corresponds to n^2 of the $\sqrt{2}(\sigma_i \otimes \sigma_i + \sigma_j \otimes \sigma_j)$ eigenvalues; as a result, $I = \text{Tr}(\mathcal{B}\rho) \sim \lambda = +2\sqrt{2}, 0, -2\sqrt{2}$ with multiplicities: 1/4, 1/2, and 1/4, respectively. Labeling states of $k = 1, \dots, n$ vectors from basis sets: $\{\widehat{a}_{A_k}\}, \{\widehat{b}_{B_k}\}, \{\widehat{a}_{A_k}^{-1}\}, \{\widehat{b}_{B_k}^{-1}\}$ in two dimensional space as $|\Phi_\pm\rangle_k$ and $|\Psi_\pm\rangle_k$, after initialization from Equation (46), we obtain:

$$|\Phi_\pm\rangle_k |0\rangle = \frac{1}{\sqrt{2}} (k|0\rangle|0\rangle_a \otimes k|0\rangle|0\rangle_b \pm k|0\rangle|1\rangle_{a^{-1}} \otimes k|0\rangle|1\rangle_{b^{-1}}), \\ |\Psi_\pm\rangle_k |0\rangle = \frac{1}{\sqrt{2}} (k|0\rangle|0\rangle_a \otimes k|0\rangle|1\rangle_{b^{-1}} \pm k|0\rangle|1\rangle_{a^{-1}} \otimes k|0\rangle|0\rangle_b), \tag{48}$$

where for $|\Psi_\pm\rangle_k$ states the corresponding eigenvalues are $\lambda_k = \pm 2\sqrt{2}$, respectively (with maximal eigenvalue being $\lambda_{k, \max} = +2\sqrt{2}$). For all $|\Phi_\pm\rangle_k$ states eigenvalues are zero, $\lambda = 0$. Consequently, the maximal violation of CHSH inequality is obtained for combinations of $|\Psi_\pm\rangle_k$ states which yield also maximal value for $I = \text{Tr}(\mathcal{B}\rho)$ [62] equivalent to the maximal eigenvalue $\lambda_{k, \max} = +2\sqrt{2}$. This result is

in agreement with Equation (44) for $p = 0$, with direct repercussion to notion of the non-local correlations [55] based on non-Abelian set of rank two free group F_2 .

6. Conclusions

A major advance towards the characterization of complexity of dynamical systems has affected communication complexity as well, offering new paths towards successful implementation of quantum information processing, especially in the field of topological insulators. In particular, the topological Rényi entropy has qualified as a good probe of the topological order, where the amount of fractal distribution present in the system and its scaling are essential for distinguishing between different phases of matter. In addition to former, wide-range implications of here presented results include the fundamental result about distribution and presence of quantum correlation and non-locality in hypergraph state based on free group on two generators, proceeding to direct implementations of underlying topology of the free group F_2 fractal sets in chip integrated circuits for quantum computing. We have defined the fractional Rényi entropy of order q for the hypergraph fractal sets based on a (non-Abelian) free group on finite generators and shown that intractability of the fractal dynamical processes can be efficiently bypassed using the geometrical concept of Lyapunov, which has proved to be the most viable method for the investigation of the complexity evolution, having its intrinsic relation to the topological and the metric (information) entropy.

Acknowledgments

The author acknowledges partial financial support through the SigmaPhy2014 organization. The author is indebted to Gavin Brennen for fruitful discussions. The work is funded through the grant: ObAd: 1007211.

Conflicts of Interest

The author declares no conflict of interest.

References

1. Gühne, O.; Cuquet, M.; Steinhoff, F.E.S.; Moroder, T.; Rossi, M.; Bruß, D.; Kraus, B.; Macchiavello, C. Entanglement and nonclassical properties of hypergraph states. *J. Phys. A* **2014**, *47*, 335303.
2. Katok, A.; Hasselblatt, B. *Introduction to the Modern Theory of Dynamical Systems*; Cambridge University Press: Cambridge, UK, 1995.
3. Hasselblatt, B.; Katok, A. Entropy and Chaos. In *A First Course in Dynamics with a Panorama of Recent Developments*; Cambridge University Press: Cambridge, UK, 2003; pp. 242–256.
4. Katok, A. Lyapunov exponents, entropy and periodic orbits for diffeomorphisms. *Publ. Math. IHES* **1980**, *51*, 137–174.
5. Oseledets, V.I.A. Multiplicative ergodic theorem: Characteristic Lyapunov exponents of dynamical systems. *Trans. Moscow Math. Soc.* **1968**, *19*, 197–231.

6. Adler, R.; Konheim, A.; McAndrew, M. Topological entropy. *Trans. Amer. Math. Soc.* **1965**, *114*, 309–319.
7. Adler, R.; Marcus, B. *Topological entropy and equivalence of dynamical systems*; Memoirs of the American Mathematical Society: Providence, RI, USA, 1979; Volume 20, pp. 219.
8. Shannon, C.E. A mathematical theory of communication. *Bell Syst. Tech. J.* **1948**, *27*, 417–436, 623–656.
9. Misiurewicz, M.; Szlenk, W. Entropy of piecewise monotone mappings. *Studia Math.* **1980**, *67*, 45–63. MR 82a:58030.
10. Landwehr, N.; Hall, M.; Frank, E. Logistic Model Trees. *Mach. Learn.* **2005**, *59*, 161–205.
11. Alsedà, L.; Llibre, J.; Misiurewicz, M. Periodic orbits of maps of Y . *Trans. Am. Math. Soc.* **1989**, 475–538.
12. Blanchard, F. A disjointness theorem involving topological entropy. *Bull. Soc. Math. France* **1993**, *121*, 465–478. MR1254749 (95e:54050).
13. Nilson, C.; Bernardes, Jr. On the set of points with a dense orbit. *Proc. Am. Math. Soc.* **2000**, *128*, 3421–3423.
14. Danilenko, A. Entropy theory from the orbital point of view. *Monatsh. Math.* **2001**, *134*, 121–141.
15. Coxeter, H.S.M.; Moser, W.O.J. *Generators and relations for discrete groups*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, 3rd ed.; Springer-Verlag: Berlin, Germany, 1972; Volume 14.
16. Kharlampovich, O.; Myasnikov, A. Elementary theory of free non-Abelian groups. *J. Algebra* **2006**, *302*, 451–552.
17. Falconer, K. *Fractal Geometry: Mathematical Foundations and Applications*; Wiley: Chichester, UK, 2004; p. 308.
18. Rényi, A. On measures of entropy and information. In Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Berkeley, CA, USA, 1960; University of California Press: Oakland, CA, USA; Volume 1, pp. 547–561.
19. Rossi, M.; Huber, M.; Bruß, D.; Macchiavello, C. Quantum hypergraph states. *New J. Phys.* **2013**, *15*, 113022.
20. Shor, P.W. Scheme for reducing decoherence in quantum computer memory. *Phys. Rev. A* **1995**, *52*, R2493–R2496.
21. Raussendorf, R.; Harrington, J. Fault-tolerant quantum computation with high threshold in two dimensions. *Phys. Rev. Lett.* **2007**, *98*, 190504.
22. Gottesman, D. Stabilizer Codes and Quantum Error Correction. Ph.D. Dissertation, California Institute of Technology, Pasadena, CA, USA, 1997.
23. Brun, T.; Devetak, I.; Hsieh, M.H. Correcting quantum errors with entanglement. *Science* **2006**, *314*, 436–439.
24. Djordjevic, I. *Quantum Information Processing and Quantum Error Correction: An Engineering Approach*; Academic Press: Oxford, UK, 2012; p. 347.
25. Berge, C. *Graphs and Hypergraphs*, 2nd ed.; North Holland: Amsterdam, The Netherlands, 1976.
26. De la Harpe, P. *Topics in Geometric Group Theory*; Chicago Lectures in Mathematics; University of Chicago Press: Chicago, IL, USA, 2000.
27. Runde, V. *Lectures on Amenability. Lecture Notes in Mathematics*; Springer-Verlag: Berlin, Germany, 2002; Volume 1774.

28. Stern, A. Non-Abelian states of matter. *Nature* **2010**, *464*, 187–193.
29. Paterson, A.L.T. *Amenability, Mathematical Surveys and Monographs*; American Mathematical Society: Providence, RI, USA, 1988; Volume 29.
30. Følner, E. On groups with full Banach mean value. *Math. Scand.* **1955**, *3*, 243–254.
31. Bollobás, B., Ed. *Advances in Graph Theory. Annals of Discrete Mathematics*; North-Holland: Amsterdam, The Netherlands, 1978; Volume 3.
32. Cayley, A. A theorem on trees. *Quart. J. Math.* **1889**, *23*, 376–378.
33. Von Neumann, J. Zur allgemeinen Theorie des Masses. *Fundam. Math.* **1929**, *13*, 73–116.
34. Mor, M.; Fraenkel, A.S. Cayley permutations. *Discrete Math.* **1984**, *48*, 101–112.
35. Lesh, N.; Marks, J.; Patrignani, M. Interactive partitioning system demonstration, short. In *Graph Drawing. Lecture Notes in Computer Science*; Springer-Verlag: Berlin, Germany, 2001; Volume 1984, pp. 31–36.
36. Brown, R.; Higgins, P.J.; Sivera, R. Nonabelian Algebraic Topology: Filtered spaces, Crossed Complexes, Cubical Homotopy Groupoids, EMS Tracts in Mathematics; European Mathematical Society: Zürich, Switzerland, 2010; Volume 15.
37. Falconer, K.J. *The Geometry of Fractal Sets*; Cambridge tracts in mathematics; University of Bangalore Press: Bangalore, India, 1997; Volume 85, p. 120.
38. Gerald, A.E. Integral, Probability, and Fractal Measures (Mathematics and Theoretical Computer); Springer-Verlag: New York, NY, USA, 1998; p. 3.
39. Stein, E.M.; Shakarchi, R. *Measure Theory, Integration, and Hilbert Spaces*; Princeton Lectures in Analysis; Princeton University Press: Princeton, NJ, USA, 2005; pp. 325–327.
40. Mandelbrot, B.B. *The Fractal Geometry of Nature*; Freeman: San Francisco, CA, USA, 1982.
41. Berge, C. *Fractional Graph Theory*; ISI Lecture Notes; Macmillan of India: New Delhi, India, 1978; Volume 1.
42. Blanchard, F., Maass, A., Nogueira, A., Eds. *Topics in Symbolic Dynamics and Applications*; Cambridge University Press/London Mathematical Society: London, UK, 2000; Volume 279.
43. Hasselblatt, B., Katok, A., Eds. *Handbook of Dynamical Systems*; Elsevier: Amsterdam, The Netherlands, 2002; Volume 1A.
44. Adler, R.L.; Coppersmith, D.; Hassner, M. Algorithms for sliding block: An application of symbolic dynamics to information theory. *IEEE Trans. Inf. Theory* **1983**, *29*, 5–22
45. Samelson, H. On the Perron-Frobenius theorem. *Michigan Math. J.* **1957**, *4*, 57–59.
46. Bronzi, M.; Tahzibi, A. Homoclinic tangency and variation of entropy. *Cad. Mat.* **2009**, *10*, 133–143.
47. Pesin, Y. Characteristic Lyapunov exponents and smooth ergodic theory. *Russ. Math. Surveys* **1977**, *32*, 55–114.
48. Kolmogorov, A.N. Entropy per unit time as a metric invariant of automorphism. *Dokl. Acad. Nauk.* **1959**, *124*, 754–755.
49. Sinai, Y.G. On the Notion of Entropy of a Dynamical System. *Dokl. Acad. Nauk.* **1959**, *124*, 768–771.
50. Kaplan, J.L.; Yorke, J.A. Chaotic Behavior of Multidimensional Difference Equations. In *Functional Differential Equations and Approximations of Fixed Points*; Lecture Notes in

- Mathematics; Peitgen, H.-O., Walter, H.-O., Eds.; Springer: Berlin, Germany, 1979; Volume 730, p. 204.
51. MacKernan, D.; Nicolis, G. Generalized Markov coarse graining and spectral decompositions of chaotic piecewise linear maps. *Phys. Rev. E* **1994**, *50*, 988.
 52. Sato, S.; Sano, M.; Sawada, Y. Practical methods of measuring the generalized dimension and the largest Lyapunov exponent in high dimensional chaotic systems. *Prog. Theor. Phys.* **1987**, *77*, 1–5.
 53. Krieger, W. On entropy and generators of measure-preserving transformations. *Trans. Amer. Math. Soc.* **1970**, *149*, 453–464.
 54. Popescu, S.; Rohrlich, D. Quantum non-locality as an axiom. *Found. Phys.* **1994**, *24*, 379–385.
 55. Brennen, G.K.; Iblisdir, S. Pachos, J.K.; Slingerland, J.K. Non-locality of non-Abelian anyons. *New J. Phys.* **2009**, *11*, 103023.
 56. Ceccherini-Silberstein, T.; Grigorchuk, R.; de la Harpe, P. Amenability and paradoxical decompositions for pseudogroups and for discrete metric spaces. *Proc. Stek. Inst. Math.* **1999**, *224*, 57–97.
 57. Wagon, S. *The Banach-Tarski Paradox*; Cambridge University Press: New York, NY, USA, 1985.
 58. Clauser, J.F.; Horne, M.A.; Shimony, A.; Holt, R.A. Proposed experiment to test local hidden-variable theories. *Phys. Rev. Lett.* **1969**, *23*, 880–884.
 59. Cirel'son, B.S. Quantum generalizations of Bell's inequality. *Lett. Math. Phys.* **1980**, *4*, 93–100.
 60. Wang, R.Q.J.; Li, Z.; Bao, Y. Encoding hypergraphs into quantum states. *Phys. Rev. A* **2013**, *87*, 022311.
 61. Berc, V. Phase space dynamics and control of the quantum particles associated to hypergraph states. **2014**, arXiv: 1411.4059.
 62. Bell, J.S. On the Einstein-Podolsky-Rosen paradox. *Physics* **1964**, *1*, 195–200.
 63. Barrett, J.; Linden, N.; Massar, S.; Pironio, S.; Popescu, S.; Roberts, D. Non-local correlations as an information-theoretic resource. *Phys. Rev. A* **2005**, *71*, 022101:1–022101:11.

© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).