





Article

Synthesis, In Silico, and In Vitro Evaluation of Anti-Leishmanial Activity of Oxadiazoles and Indolizine Containing Compounds Flagged against Anti-Targets

Strahinja Stevanovic¹, Milan Sencanski¹, Mathieu Danel² , Christophe Menendez^{3,4}, Roumaissa Belguedj^{3,4,5}, Abdelmalek Bouraiou⁵, Katarina Nikolic⁶ , Sandrine Cojean⁷, Philippe M. Loiseau^{7,*}, Sanja Glisic^{1,*} , Michel Baltas^{3,4,*} and Alfonso T. García-Sosa^{8,*} 

¹ Laboratory for Bioinformatics and Computational Chemistry, Institute of Nuclear Sciences VINCA, University of Belgrade, P.O. Box 522, 11001 Belgrade, Serbia; strahinja.stevanovic@protonmail.com (S.S.); sencanski@vin.bg.ac.rs (M.S.)

² ITAV, Université de Toulouse, CNRS, 31062 Toulouse, France; mathieu.danel@itav.fr

³ Department of Chemistry, Université de Toulouse, UPS, CNRS UMR 5068, LSPCMIB, 118 Route de Narbonne, 31062 Toulouse, France; menendez@chimie.ups-tlse.fr (C.M.); racha_belg@hotmail.fr (R.B.)

⁴ CNRS, Laboratoire de Synthèse et Physico-Chimie de Molécules d'Intérêt Biologique, LSPCMIB, UMR-5068, 118 Route de Narbonne, 31062 Toulouse, France

⁵ Unité de Recherche de Chimie de l'Environnement et Moléculaire Structurale, Université Frères Mentouri, Route de Ain El Bey, 25000 Constantine, Algeria; bouraiou.abdelmalek@yahoo.fr

⁶ Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Belgrade, Vojvode Stepe 450, 11000 Belgrade, Serbia; knikolic@pharmacy.bg.ac.rs

⁷ Antiparasitic Chemotherapy, UMR 8076 CNRS BioCIS, Faculty of Pharmacy Université Paris-Sud, Rue Jean-Baptiste Clément, F 92290 Chatenay-Malabry, France; sandrine.cojean@u-psud.fr

⁸ Institute of Chemistry, University of Tartu, Ravila 14a, 50411 Tartu, Estonia

* Correspondence: philippe.loiseau@u-psud.fr (P.M.L.); sanja@vin.bg.ac.rs (S.G.); baltas@chimie.ups-tlse.fr (M.B.); alfonsog@ut.ee (A.T.G.-S.)

Academic Editor: Thomas J. Schmidt

Received: 26 February 2019; Accepted: 28 March 2019; Published: 2 April 2019



Abstract: Due to the lack of approved vaccines against human leishmaniasis and the limitations of the current chemotherapy inducing side effects and drug resistance, development of new, effective chemotherapeutic agents is essential. This study describes the synthesis of a series of novel oxadiazoles and indolizine-containing compounds. The compounds were screened in silico using an EIIP/AQVN filter followed by ligand-based virtual screening and molecular docking to parasite arginase. Top hits were further screened versus human arginase and finally against an anti-target battery to tag their possible interactions with proteins essential for the metabolism and clearance of many substances. Eight candidate compounds were selected for further experimental testing. The results show measurable in vitro anti-leishmanial activity for three compounds. One compound with an IC₅₀ value of 2.18 μ M on *Leishmania donovani* intramacrophage amastigotes is clearly better positioned than the others as an interesting molecular template for further development of new anti-leishmanial agents.

Keywords: Leishmania; arginase; in silico; anti-target; in vitro; anti-leishmanial inhibitors; anti-target

1. Introduction

The leishmaniasis are a group of diseases caused by the trypanosomatid protozoan parasite of the genus *Leishmania* through the bite of infected phlebotomine sandflies, endemic to 97 countries in

parts of the tropics, subtropics, and in Southern Europe [1]. An estimated 700,000 to 1 million new cases of leishmaniasis and 20,000–30,000 deaths occur annually. In 2017, 94% of new cases reported to the WHO occurred in seven countries: Brazil, Ethiopia, India, Kenya, Somalia, South Sudan, and Sudan [2]. The three main clinical syndromes of leishmaniasis include cutaneous leishmaniasis (CL), muco-cutaneous leishmaniasis (MCL), and visceral leishmaniasis (VL) [3].

Leishmaniasis is one of the most neglected diseases in terms of drug development [4,5]. Without any effective vaccine against leishmaniasis in humans at present, the control of these parasites relies solely on chemotherapy [6]. The current chemotherapies suffer from several drawbacks due to toxicity, long-term treatment, severe adverse effects, high cost, invasive administration routes, low effectiveness, and commonly, drug resistance [4,7,8]. Therefore, there is an urgent need for novel compounds with substantial antileishmanial effect and low toxicity to the host.

Leishmania and all members of the *Trypanosomatidae* family depend on polyamines (PA) for growth and survival [9,10]. The first enzyme involved in PA biosynthesis that hydrolyses arginine into ornithine and urea is arginase [11]. Arginase inhibition can lead to oxidative stress in parasite cells owing to a deficiency in trypanothione production that neutralizes reactive macrophage-derived oxygen and nitrogen species [12], consequently leading to infection control [13]. Arginase is important for parasite survival: deleting the arginase gene is auxotrophic for polyamines [14]. Being arginase distinct from the mammalian target [11] and absolute necessity of the enzyme for survival of the pathogen, it represents a key parasite drug target [15].


Oxadiazoles have been reported as potent anti-leishmanial agents [16,17]. Potent antimicrobial and anti-leishmanial activity are also documented for indolizine-containing compounds [18,19]. The MetIDB database was screened in previous work with *in silico* procedures and ten promising flavonoids were proposed as anti-leishmanial candidates [20]. In this study, a series of novel oxadiazoles and indolizine containing compounds were synthesized, screened *in silico* with the Electron Ion Interaction Potential/Average Quasi Valence Number (EIIP/AQVN) filter, followed by ligand-based virtual screening and molecular docking to *Leishmania* arginase. In addition, the top hits were virtually screened against human arginase and anti-targets. The best selected candidates for anti-leishmanial compounds were subjected to experimental testing. The experimental results show measurable *in vitro* anti-leishmanial activity for three compounds.

2. Results

2.1. Chemistry











2.1.1. Synthesis of 1,2,4-Oxadiazoles

N-acylhydrazones are important and versatile scaffolds. They can be obtained by conventional [21] or non-conventional methods [22,23]. They can be considered either as final products with important biological properties or stable intermediates in the design and construction of other motifs like 1,2,4-oxadiazoles. In our hands, as *N*-acylhydrazones 1–5 were prepared on an automated platform, we conducted synthesis by refluxing an equimolar mixture of hydrazide and aldehyde in ethanol with a catalytic amount of hydrochloric acid (Scheme 1). Purifications by filtration of the solid compounds were parallelized and we were able to isolate about 200 compounds in four batches. The five compounds used in this work are shown in Table 1.



Scheme 1. Automated synthesis of acylhydrazones 1–5; for compounds that were isolated: R₁ = furan, thiophene, pyridine; R₂ = 4-methyl-thiazole, 2-bromo-thiophene, benzothiazole.


Table 1. Isolated acylhydrazones.

Compound	Hydrazide	Aldehyde	Yield (%)	Purity ¹ (%)
1			61	92
2			72	99
3			71	99
4			quant.	99
5			70	92

¹ Determined by Liquid Chromatography/Mass Spectrometry (LC/MS).

1,2,4-Oxadiazoles can be considered as one of the most important 5-membered heteroaromatic rings found in many pharmaceutical compounds. Among the various synthetic approaches reported in the literature [24], one concerns reaction under conventional or non-conventional methods of amidoximes with suitably activated acid derivatives [25].

Synthesis of the 3-[3,4-(methylenedioxy)phenyl]-5-(methoxymethyl)-1,2,4-oxadiazole **8** was achieved in two steps and 90% total yield by the reaction of 3,4-(methylenedioxy)benzonitrile with hydroxylamine to afford quantitatively amidoxime **6** followed by the reaction with methoxyacetyl chloride under microwave irradiation (Scheme 2). This compound was prepared during a small molecule library synthesis program, and the protocols established to allow the automation and the parallelization of reactions. The first step of the procedure was applied to 3-nitrobenzonitrile, affording also in quantitative yield the corresponding 3-nitrobenzamidoxime.




Scheme 2. Automated synthesis of amidoximes **6**, **7** and 1,2,4-oxadiazole **8**.


2.1.2. Synthesis of Indolizines

The 1,3-dipolar cycloaddition reaction of azomethine ylide containing nitrogen in a 6-membered ring such as pyridine and isoquinoline was applied for the synthesis of the target compounds. In that

respect, 1-((5-nitro-1*H*-benzo[d]imidazol-2-yl)methyl)pyridin-1-ium chloride **9** and 2-((5-nitro-1*H*-benzo[d]imidazol-2-yl)methyl)isoquinolin-2-ium chloride **10**(2), (Scheme 3), which were prepared from the reaction of pyridine/isoquinoline with 2-(chloromethyl)-5-nitro-1*H*- benzo[d]imidazole (Scheme 3), as described previously [26,27], undergo a dehydrohalogenation reaction in the presence of triethylamine in chloroform to give the corresponding pyridinium **9**/isoquinolinium **10** ylide in 75% and 95% yield, respectively. The reaction of the latter (Scheme 4) with DMAD or methyl propiolate at room temperature and overnight afforded, after workup and column chromatography, the indolizine **11** (25% yield), and benzindolizines **12** and **13** (28% and 27%, respectively). The obtained compounds were fully characterized by ^1H , ^{13}C NMR, mass spectra, and IR.



Scheme 3. Preparation of 1-((5-nitro-1*H*-benzo[d]imidazol-2-yl) methyl) pyridin-1-ium chloride **9** and 2-((5-nitro-1*H*-benzo[d]imidazol-2-yl) methyl) isoquinolin-2-ium chloride **10**.



Scheme 4. Formation of indolizine derivatives **11–13**.

2.2. Virtual Screening

2.2.1. EIIP Filtering

We used the previously developed average quasi valence number/electron ion interaction potential (EIIP/AQVN) criteria for selection of *Leishmania* arginase compounds [20].

2.2.2. 3D QSAR Filtering

39 molecules were obtained after filtering with EIIP criteria and were subjected to prediction of their activity using the above described arginase 3D-quantitative structure-activity relationship (3D-QSAR) model. The two criteria for selection were: (1) partial least square (PLS) scores in the

vicinity of compounds from the model (Figure 1); and (2) best ranking by predicted IC_{50} values. This filtering gave 10 candidates that were used for docking into the *Leishmania* arginase structure model, human arginase crystal structure, and off-target (anti-target) affinity calculation.




Figure 1. Partial least squares (PLS) scores of 39 candidate molecules (red dots) and training set (black dots). The best candidate compound, 2, is marked (PICA052).

2.2.3. Arginase Docking

Ten compounds were docked into both the parasite arginase model structure and the crystal structure of human arginase. Docking scores of the best-docked conformations, along with experimental measurements are presented in Table 2. Although PLS prediction and docking score results were promising, the experiments showed significant activity of six compounds. However, due to their toxicity, only one candidate, compound 2, is acceptable thanks to its selectivity index (SI, CC_{50} for macrophage/ IC_{50} for promastigotes) ≥ 2 , with an IC_{50} value on intramacrophage amastigotes $< 5 \mu\text{M}$. The highest ranked docking conformation of this compound in the *Leishmania* arginase model is presented in Figure 2.












Figure 2. Best candidate compound 2, docked into the active site of *Leishmania* arginase, with marked aminoacids and important interactions. Green: hydrogen bond, orange: electrostatic interactions, purple: hydrophobic interactions.

Table 2. Compounds selected from QSAR PLS scores, with calculated EIIP descriptor values, predicted activity, docking scores, and experimental data.

No	Code	AQVN	EIIP	Structure	Pred. pIC ₅₀	Autodock Score (kcal/mol) <i>Leishmania Arginase</i>	Autodock Score (kcal/mol) <i>Human Arginase</i>	<i>Leishmania donovani</i> LV9 axenic Amastigotes Forms IC ₅₀ ± SD (μM)	<i>Leishmania donovani</i> Intramacrophage Amastigotes Forms IC ₅₀ ± SD (μM)	Toxicity on RAW 264.7 Macrophages CC ₅₀ ± SD (μM)
17	1	3.28	0.125		3.85	-6	-5.9	11.89 ± 1.87	6.25 ± 1.89	13.86 ± 2.11
21	2	3.304	0.129		3.77	-5.8	-5.0	1.09 ± 0.12	2.18 ± 0.01	4.87 ± 0.39
22	3	3.304	0.129		3.94	-6.2	-5.8	> 100	> 100	> 100
23	4	3.357	0.134		3.55	-6.3	-6.2	12.50 ± 0.61	11.22 ± 0.62	16.34 ± 1.42
24	5	3.267	0.123		3.54	-6.4	-6.3	61.52 ± 5.99	55.22 ± 4.77	>100
25	8	3.259	0.122		3.77	-7.4	-6.8	59.86 ± 6.45	51.26 ± 4.35	>100
26	7	3.4	0.134		5.39	-7.4	-7.3	51.21 ± 4.98	47.24 ± 3.12	>100
38	12	3.3	0.128		3.94	-7.6	-6.8	>100	>100	>100

2.2.4. Anti-Target Interaction Matrix

The top eight compounds from the filtering were then assessed against the anti-target battery. The results of the docking of all the final compounds against the battery of five anti-targets are shown in Figure 3 (full table of docking scores Table 3).

	PXR	SULT	CYP 2a6	CYP 2c9	CYP 3a4	Total
2						0
3						1
4						0
5						1
1						0
8						0
7						0
12						0.5

Figure 3. Interaction matrix between proposed ligands and anti-targets. Color code: black = 1.0; grey = 0.5, white = 0.0. Columns: PXR = pregnane-X-receptor; SULT = sulfotransferase; CYP 2a6 = cytochrome P450 cam 2a5; CYP 2c9 = cytochrome P450 cam 2c9; CYP 3a4 = cytochrome P450 cam 3a4; Total = PXR + SULT + CYP 2a6 + CYP 2c9 + CYP 3a4.

Table 3. In vitro anti-leishmanial activity and cytotoxicity of compounds selected by virtual screening.

Compound	<i>Leishmania donovani</i> LV9 Axenic Amastigotes IC ₅₀ ± SD (µM)	<i>Leishmania donovani</i> LV9 Intramacrophage Amastigotes IC ₅₀ ± SD (µM)	Cytotoxicity on RAW 264.7 Macrophages CC ₅₀ ± SD (µM)	SI = CC ₅₀ /IC ₅₀ Intramacro. Amas.
2	1.09 ± 0.12	2.18 ± 0.12	4.87 ± 0.39	2.2
3	>100	>100	>100	/
4	12.50 ± 0.62	11.22 ± 0.62	16.34 ± 1.42	1.4
5	61.52 ± 5.99	55.22 ± 4.77	> 100	>1.8
1	11.89 ± 1.87	6.25 ± 1.89	13.86 ± 2.11	2.2
8	59.86 ± 6.45	51.26 ± 4.35	>100	>1.9
7	51.21 ± 4.98	47.24 ± 3.12	>100	>2.1
12	>100	>100	>100	/
AmB	0.20 ± 0.08	0.11 ± 0.09	5.36 ± 0.52	48.7

There was broad general agreement between the five anti-targets. None of the compound's docking score surpassed that of the threshold for CYP P450 2a6, 2c9, or 3a4, which may be an indication of the relative size of the ligands. There were more interactions found with the anti-targets PXR and SULT. Compounds 3 and 5 had the highest combined score of 1.0 for all anti-targets combined, while most compounds had even lower interactions.

All of the compounds had zero PAINS flags, passed Lipinski's rule-of-five, and were predicted to be soluble or moderately soluble, have high gastrointestinal absorption, and potential (using a support-vector-machine) CYP 2a6 binding for 2, 3, 4, 5, and CYP 3a4 for 8, according to filters [28].

2.2.5. In vitro Evaluation of Anti-Leishmanial Activity

Finally, we selected eight hit compounds for experimental testing. Since isolated arginase was not available, the selected compounds were assayed for their in vitro inhibition activity against axenic amastigote and intramacrophage amastigote forms of *Leishmania donovani* (Table 3). A broad range of activities against axenic and intramacrophage amastigotes forms of *Leishmania donovani* was found with IC₅₀ values in the range between 1–2 and 55–61 µM. The compounds exhibited similar activities on both axenic amastigotes and intramacrophage amastigotes. The most potent inhibitors were also slightly toxic with CC₅₀ values in the range from 4 to 16 µM.

Regarding structure–activity relationships, the replacement of a thiophene group (compound 4) by a pyridine group (compound 5) was responsible for a loss of activity by a factor of five and also a strong reduction of cytotoxicity. Among the tested compounds, one compound, 2, is the most active despite having a not negligible toxicity. Its chemical structure has two thiophene groups. The replacement of a thiophene group (compound 2) by a furan group (compound 3) led to a total loss of activity and cytotoxicity. The replacement of the bromothiophene group (compound 2) by a benzothiazole group (compound 4) was responsible for a five-fold reduction of the anti-leishmanial activity and four-fold reduction of cytotoxicity. The replacement of bromothiophene (compound 3) by a methylbenzothiazole group (compound 1) enhanced both the antileishmanial activity and cytotoxicity. In the phenotypic based screening, the most promising compound 2 (Table 3) has an IC_{50} value of 2.18 μ M, 20 times higher than that of amphotericin B, the reference compound. Regarding the selectivity index (SI) values, compound 2 has an SI of 2, whereas amphotericin B had an SI of 48 against *Leishmania axenic amastigotes*. It is also less than four times higher than the reference drug miltefosine (0.31 μ M) [29]. The obtained in vitro results clearly require additional pharmacomodulations to reduce the cytotoxicity and enhance the antileishmanial activity in order to achieve better selectivity index values.

3. Discussion

Inhibition of enzymes of the polyamine–trypanothione metabolism including arginase is considered as one of the best options for the treatment of *Leishmania* since many of these enzymes passed both target validation and chemical validation [10]. In the quest for potential treatment options against leishmaniasis, increasing interest has been shown for *N*-acylhydrazones, oxadiazole, and indolizine containing compounds. These types of compounds targeting arginase represent an interesting strategy in the search for a new anti-leishmanial treatment.

In our hands, *N*-acylhydrazones were prepared on an automated platform where reaction conditions and purifications by filtration of the solid compounds were parallelized for 50 compounds in each run (about 200 compounds synthesized in four batches). In that respect, reaction conditions were generalized, and for some of the compounds, yields are not optimum. The same is also true for the 1,2,4-oxadiazoles where the reaction conditions were chosen in order to prepare a small-molecule library (150 compounds synthesized). The reaction conditions first afforded the amidoxime compounds quantitatively, then the reaction with activated acid furnished in good yield the 1,2,4 oxadiazoles, where one example is used in the current study. Concerning the indolizine compounds, the azomethine ylide prepared in situ by 1,3 dipolar cycloaddition was chosen. The yields obtained for compounds 11–13 are poor, varying between 25–28% after careful purification. The insolubility of the chloride salts could be the reason for the low yields. As one of the research programs of the group concerns 1,3 dipolar cycloadditions using this type of ylides, efforts are currently oriented to the synthesis of indolizine-bearing derivatives under non-conventional methods in order to obtain focused small libraries of these compounds.

The compounds thus obtained for this study were subsequently screened. The ten selected compounds from our study with favorable in silico interaction profiles with arginase and against the anti-targets were favorable for further experimental testing and represent better initial points than screening compounds at random. Three compounds were found to be active against *Leishmania donovani*. One product, compound 2, is better than the others as an interesting molecular template for further development of new anti-leishmanial agents due to its observed experimental activity against parasite amastigotes, better selectivity index, and predicted interactions with targets and anti-targets. Interestingly, the compound that had the best selectivity index, the most promising compound 2, had a docking score that was approximately 0.8 kcal/mol stronger for the parasite arginase than for human arginase. This may also indicate that a larger degree of selectivity for a compound may be picked up by the procedure carried out here.

Considering the mechanism of action of both chemical series, the results obtained in this paper prompt us to develop further pharmacomodulations in order to diminish the cytotoxicity and enhance

the anti-leishmanial activity. In any case, despite the level of cytotoxicity of compound **2** being in a similar range to that of the reference compound AmB, the obtained results justify the determination of the maximal tolerated dose in mice, and then, the in vivo evaluation of compound **2** on the *L. donovani*/BALB/c mice model.

4. Materials and Methods

4.1. Chemistry and Physico-Chemical Analyses


Melting points (m.p.) were determined using a Mettler Toledo MP50 system and were uncorrected. ^1H and ^{13}C NMR spectra were recorded in CDCl_3 and/or DMSO-d_6 using a Bruker AC 300 (^1H) or 75 MHz (^{13}C) instruments, except for compounds **1**, **2**, **4**, **6**, and **7**: ^{13}C -NMR analyses were conducted with a Bruker AVANCE500 at 298 K (125.75 MHz). Chemical shifts are given in δ parts per million (ppm) and referenced to external TMS.

Automated syntheses were carried out on an Accelerator SLT-106 workstation from Chemspeed and microwave assisted reactions on a SWave workstation from Chemspeed equipped with a Biotage Initiator reactor.


High-resolution mass spectra (MS) were recorded on a GCT Premier Mass Spectrometer (Waters, MA, USA). Liquid Chromatography/Mass Spectrometry (LC/MS) analyses were performed on an Autopurif system from Waters (PDA 2996, MS 3100, Pump 2545, MA, USA) with a Gemini-NX column (5, C18, 110A, 50×4.6 mm) from Phenomenex. Analyses were done with 1 mL/min flow and a gradient of water/acetonitrile containing 0.05% of formic acid (0.0 to 1.0 min: 90/10; 1.0 to 5.0 min: 90/10 to 0/100; 5.0 to 6.5 min: 0/100; 6.5 to 7.0 min: 0/100 to 90/10; 7.0 to 12.0: 90/10) and purities were determined at 254 nm.

4.1.1. General Procedure for the Automated Synthesis of Compounds **1**–**5**:

Aldehyde (1.00 mmol), hydrazide (1.00 mmol), ethanol (4.0 mL) and 50 μL of aqueous 1M HCl were added in a 13 mL double jacket reactor from Chemspeed equipped with condenser. The mixture was shaken at 700 rpm for 4 h at 80 $^\circ\text{C}$, then cooled down to 20 $^\circ\text{C}$. The precipitate was filtrated and successively washed with 2 mL of ethanol and 2 mL of ethyl ether to obtain the product.




N'-[(4-Methyl-2-thiazolyl)methylene]furan-2-carbohydrazide (**1**): Tan solid, yield: 61%, m.p.: 121–124 $^\circ\text{C}$. LC purity (254 nm): 92%. ^1H -NMR (300 MHz, DMSO) δ ppm: 12.22 (s, 1H), 8.62 (s, 1H), 8.00–7.93 (m, 1H), 7.41–7.33 (m, 2H), 6.73 (dd, $J = 3.3, 1.5$ Hz, 1H), 2.39 (d, $J = 0.9$ Hz, 3H). ^{13}C -NMR (125.75 MHz, DMSO) δ ppm: 163.72, 157.80, 154.60, 153.93, 142.57, 116.86, 116.21, 115.16, 112.74, 112.38, 17.14. HRMS (DCI- CH_4 , TOF) m/z : $[\text{M} + \text{H}]^+$ calc. for $\text{C}_{10}\text{H}_{10}\text{N}_3\text{O}_2\text{S}$: 236.0494. Found: 236.0494.




N'-[(5-Bromo-2-thiophenyl)methylene]thiophene-2-carbohydrazide (**2**): Light yellow solid, yield: 72%, m.p.: 195–198 $^\circ\text{C}$. LC purity (254 nm): 99%. ^1H -NMR (300 MHz, DMSO) δ ppm: 11.89 (s, 1H), 8.58 and 8.20 (2 broad singlets, 1H), 8.05–7.93 (m, 1 H), 7.88 (br, 1H), 7.33 (dd, $J = 4.5, 0.6$ Hz, 1H), 7.27 (d, $J = 3.9$ Hz, 1H), 7.22 (dd, $J = 5.1, 3.9$ Hz, 1H). ^{13}C -NMR (125.75 MHz, DMSO) δ ppm: 161.58, 158.40, 142.42, 141.41,


141.24, 138.50, 138.15, 135.65, 135.26, 133.46, 132.47, 131.98, 129.51, 128.60, 127.14, 115.25. HRMS (ES, TOF) m/z : $[M + H]^+$ calc. for $C_{10}H_8N_2OS_2Br$: 314.9261. Found: 314.9250.



N'-[(5-Bromo-2-thiophenyl)methylene]furan-2-carbohydrazide (**3**): Light yellow solid, yield: 71%, m.p.: 201–202 °C. LC purity (254 nm): 99%. 1H -NMR (300 MHz, DMSO) δ ppm: 11.89 (s, 1H), 8.57 (br, 1H), 7.94 (s, 1H), 7.32–7.24 (m, 3H), 6.70 (s, 1H). ^{13}C -NMR (75 MHz, DMSO) δ ppm: 154.1, 146.5, 146.0, 142.1, 140.9, 131.5, 131.3, 115.1, 114.8, 112.1. HRMS (ES, TOF) m/z : $[M + H]^+$ calc. for $C_{10}H_8N_2O_2SBr$: 298.9490. Found: 298.9492.




N'-[(2-Benzothiazolyl)methylene]thiophene-2-carbohydrazide (**4**): Light yellow solid, yield: quant., m.p.: 231–234 °C. LC purity (254 nm): 99%. 1H -NMR (300 MHz, DMSO) δ ppm: 12.37 (s, 1H), 8.74 and 8.42 (2 br, 1H), 8.20–8.12 (m, 1H), 8.12–7.87 (m, 3H), 7.55 (ddd, $J = 14.7, 7.5, 1.5$ Hz, 1H), 7.50 (ddd, $J = 15.9, 7.5, 1.5$ Hz, 1H), 7.27 (dd, $J = 4.8, 3.6$ Hz, 1H). ^{13}C -NMR (125.75 MHz, DMSO) δ ppm: 165.52, 164.86, 161.86, 158.45, 153.62, 142.24, 138.56, 135.87, 134.63, 130.34, 128.77, 127.17, 127.04, 123.76, 123.00. HRMS (ES, TOF) m/z : $[M + H]^+$ calc. for $C_{13}H_{10}N_3OS_2$: 288.0265. Found: 288.0277.




N'-[(2-Benzothiazolyl)methylene]pyridine-4-carbohydrazide (**5**): Off-white solid, yield: 70%, m.p.: 234–237 °C. LC purity (254 nm): 92%. 1H -NMR (300 MHz, DMSO) δ ppm: 12.64 (s, 1H), 8.88–8.81 (m, 2H), 8.79 (s, 1H), 8.21–8.11 (m, 1H), 8.11–8.03 (m, 1H), 7.92–7.82 (m, 2H), 7.61–7.47 (m, 2H). ^{13}C -NMR (75 MHz, DMSO) δ ppm: 164.7, 162.0, 153.1, 150.3, 143.3, 140.1, 134.1, 126.8, 126.7, 123.4, 122.6, 121.7. HRMS (ES, TOF) m/z : $[M + H]^+$ calc. for $C_{14}H_{11}N_4OS$: 283.0654. Found: 283.0656.

4.1.2. General Procedure for the Automated Synthesis of Compounds **6** and **7**:

Nitrile (20.00 mmol), ethanol (30.0 mL) and hydroxylamine (50% in water, 2.0 mL) were added in a 75 mL double jacket reactor from Chemspeed equipped with condenser. The mixture was shaken at 600 rpm for 5 h at 80 °C, then cooled down to 20 °C. The reaction medium was evaporated under reduced pressure and the solid was finally dried overnight in a heated (40 °C) vacuum oven.




3,4-(Methylenedioxy)benzamidoxime (**6**): White solid, yield: quant. 1H -NMR (300 MHz, DMSO) δ ppm: 9.51 (s, 1H), 7.22–7.16 (m, 2H), 6.93–6.88 (m, 1H), 6.03 (s, 2H), 5.72 (s, 2H). ^{13}C -NMR (125.75 MHz, DMSO) δ ppm: 150.98, 148.25, 147.56, 127.87, 119.75, 108.30, 106.17, 101.61.



3-Nitrobenzamidoxime (7): Yellow solid, yield: quant., m.p.: 182–184 °C. $^1\text{H-NMR}$ (300 MHz, DMSO) δ ppm: 9.97 (s, 1H), 8.51 (m, $J = 2.1$ Hz, 1H), 8.23 (ddd, $J = 8.1, 2.4, 0.9$ Hz, 1H), 8.12 (ddd, $J = 7.8, 1.5, 1.2$ Hz, 1H), 7.68 (t, $J = 8.1$ Hz, 1H), 6.10 (s, 2H). $^{13}\text{C-NMR}$ (125.75 MHz, DMSO) δ ppm: 149.60, 148.22, 135.40, 132.04, 130.24, 123.93, 120.40. HRMS (DCI- CH_4 , TOF) m/z : $[\text{M} + \text{H}]^+$ calc. for $\text{C}_7\text{H}_8\text{N}_3\text{O}_3$: 182.0566. Found: 182.0559.

4.1.3. Procedure for the Synthesis of Compound 8:

A mixture of 3,4-(methylenedioxy)benzamidoxime (**6**, 180 mg, 1.00 mmol), acetonitrile (3.0 mL), thiethylamine (153 μL , 1.10 mmol) and methoxyacetyl chloride (91 μL , 1.00 mmol) was placed in a sealed vial and exposed to microwave irradiation (130 °C, 1 h). The reaction medium was evaporated under reduced pressure and the solid was washed with 15 mL of ethyl acetate to give **8** (188 mg, 80% yield).




3-[3,4-(Methylenedioxy)phenyl]-5-(methoxymethyl)-1,2,4-oxadiazole (8): Light tan solid, m.p.: 56–57 °C. LC purity (254 nm): 98%. $^1\text{H-NMR}$ (300 MHz, DMSO) δ ppm: 7.58 (dd, $J = 8.1, 1.8$ Hz, 1H), 7.45 (dd, $J = 1.5, 0.3$ Hz, 1H), 7.09 (dd, $J = 8.1, 0.3$ Hz, 1H), 6.14 (s, 2H), 4.80 (s, 2H), 3.42 (s, 3H). $^{13}\text{C-NMR}$ (75 MHz, DMSO) δ ppm: 176.5, 167.2, 150.1, 148.0, 122.1, 119.6, 109.0, 106.6, 101.9, 64.4, 58.8. HRMS (DCI- CH_4 , TOF) m/z : $[\text{M} + \text{H}]^+$ calc. for $\text{C}_{11}\text{H}_{11}\text{N}_2\text{O}_4$: 235.0719. Found: 235.0712.

4.1.4. General Procedure for the Synthesis of Compounds 11–13:


Compounds 1-((5-nitro-1H-benzo[d]imidazol-2-yl)methyl)pyridin-1-ium chloride **9** and 2-((5-nitro-1H-benzo[d]imidazol-2-yl)methyl)isoquinolin-2-ium chloride **10** were prepared as previously reported and gave the same experimental characteristics. A mixture of pyridinium/isoquinolinium salt (1 mmol, 1.0 eq.) and alkyne (1.1 mmol, 1.1 eq.) in chloroform was stirred at 0 °C. Triethylamine (1.3 mmol, 1.3 eq.) was added dropwise and the mixture was further stirred at room temperature for 24 h. The reaction medium was evaporated under reduced pressure and the obtained residue was purified by flash chromatography.

99 mg of compound **11** were obtained (25% yield).




Dimethyl 3-(5-nitro-1H-benzo[d]imidazol-2-yl)indolizine-1,2-dicarboxylate (11): m.p.: 234 °C. R_f : 0.6 EP/EtOAc (7:3 v/v). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ ppm: 11.65 (s, 1H), 10.30 (d, $J = 7.2$ Hz, H5), 8.72 (d, $J = 2.1$ Hz, H4'), 8.32–8.18 (m, H6' + H8), 7.59 (d, $J = 8.8$ Hz, H7'), 7.38 (dd, $J = 9.0, 6.7$ Hz, H7), 7.11 (dd, $J = 7.2, 6.7$ Hz, H6), 4.08 (s, 3CH₃), 4.00 (s, 3CH₃). $^{13}\text{C-NMR}$ (75 MHz, DMSO) δ ppm: 169.2, 163.7, 148.2, 147.5, 147.3, 143.9, 142.6, 137.0, 173.0, 131.9, 128.4, 126.1, 119.3, 119.0, 115.2, 110.9, 107.9, 53.6, 51.8. HRMS (ES, TOF) m/z : $[\text{M} + \text{H}]^+$ calc. for $\text{C}_{19}\text{H}_{14}\text{N}_4\text{O}_6$: 395.0992. Found: 395.0988. IR (ATR) (cm^{-1}): 3347, 3117, 2954, 1716, 1702, 1503, 1215, 749.

125 mg of compound **12** were obtained (28% yield).



Dimethyl 3-(5-nitro-1H-benzo[d]imidazol-2-yl)pyrrolo[2,1-a]isoquinoline-1,2-dicarboxylate (12): m.p.: 162 °C. R_f : 0.6 EP/THF (1:4 *v/v*). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ ppm: 11.75 (s, NH), 8.98 (d, $J = 7.5$ Hz, H5), 8.59 (s, H4'), 8.35–8.26 (m, H10), 8.17 (d, $J = 8.9$ Hz, H6'), 7.86–7.83 (m, H7+H7'), 7.65–7.62 (m, H8+H9), 7.33 (d, $J = 7.5$ Hz, H6), 4.00 (s, CH_3), 3.85 (s, CH_3). $^{13}\text{C-NMR}$ (75 MHz, DMSO) δ ppm: 166.9 (CO), 164.1 (CO), 146.9 (C5'), 143.4 (C2), 129.0 (CH8 + CH9), 128.9 (C11), 128.7 (C), 128.2 (CH7), 123.9 (C), 123.9 (CH10), 123.8 (CH5), 119.3 (C), 118.7 (CH6'), 117.5 (C3), 115.4 (CH7'), 115.3 (CH6), 113.1 (CH4'), 111.0 (C), 53.3 (CH_3), 53.0 (CH_3). HRMS (ES, TOF) m/z : $[\text{M} + \text{H}]^+$ calc. for $\text{C}_{23}\text{H}_{17}\text{N}_4\text{O}_6$: 445.1148. Found: 445.1127. IR (ATR) (cm^{-1}): 3399, 3109, 2990, 1727, 1686, 1519, 1340, 1210, 734.

195 mg of compound **13** were obtained (27%)



Methyl 3-(5-nitro-1H-benzo[d]imidazol-2-yl)pyrrolo[2,1-a]isoquinoline-1-carboxylate (13): m.p.: 280 °C. R_f : 0.4 EP/EtOAc (4:1 *v/v*). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ ppm: 13.60 (s, NH), 10.03 (d, $J = 7.5$ Hz, H5), 9.78–9.73 (m, H4'), 8.54–8.51 (m, H10), 8.23 (s, H2), 8.14–8.08 (m, H6'), 7.93–7.89 (m, H7'), 7.70–7.63 (m, H7,H8,H9), 7.52 (d, $J = 7.5$ Hz, H6), 3.93 (s, CH_3). $^{13}\text{C-NMR}$ (75 MHz, DMSO) δ ppm: 164.7, 143.1, 139.5, 131.7, 129.5, 129.2, 128.0, 127.5, 127.2, 127.0, 126.1, 125.2, 124.7, 124.5, 120.3, 119.0, 116.4, 115.4, 115.1, 52.1. HRMS (ES, TOF) m/z : $[\text{M} + \text{H}]^+$ calc. for $\text{C}_{21}\text{H}_{24}\text{N}_4\text{O}_4$: 387.1067. Found: 387.1070. IR (ATR) (cm^{-1}): 2920, 2851, 1708, 1521, 1341, 1202, 760.

4.2. Ligands

The source for data was internal libraries of synthesized oxadiazoles and indolizine-containing compounds.

4.3. Virtual Screening

Starting with a filter based on EIIP/AQVN values and subsequently, 3D QSAR filtering, and arginase docking, hit compounds were docked into anti-targets involved in the metabolism of compounds in order to tag their possible interactions.

4.3.1. EIIP/AQVN

AQVN and the EIIP can give indication on long-range biomolecule interaction (over 0.5 nm) [30], based on the general model pseudopotential [31]

$$\text{EIIP} = 0.25 Z^* \sin(1.04 \pi Z^*) / 2\pi, \quad (1)$$

where Z^* is the average quasi-valence number (AQVN):

$$Z^* = \sum m(n_i Z_i / N), \quad (2)$$

where Z_i is the valence number of the i th atomic component, n_i is the number of atoms of the i th component, m is the number of atomic components in the molecule, and N is the total number of atoms. EIIP values are calculated according to equations 1 and 2 are expressed in Rydberg units (Ry).

Similarity in AQVN and EIIP values may give a clue on a common therapeutic target, thus setting up criteria for virtual screening of molecular libraries for compounds with similar therapeutic properties [20].

4.3.2. 3D QSAR

We used our previously reported 3D QSAR model of *Leishmania amazonensis* [20].

4.4. Arginase

The crystal structure of *Leishmania mexicana* with ABH (4iu0.pdb) was used to generate a model for *L. amazonensis*. Docking was performed with Autodock 4.6.2. The Lamarckian GA method was used to perform the conformational search.

4.5. Anti-Targets

The docking for the anti-targets was carried out with Glide XP [32] and proteins with resolution ≤ 2.6 Å (1m13, 2a3r, 1z10, 1og5, 1tqn) [33,34]. The thresholds for determining a strong interaction were set to -7.7 , -6.3 , -7.6 , -8.7 , and -7.5 kcal/mol. An estimate of the binding was changed to interaction codes:

$$\text{Score} = 0.0 \text{ if } \Delta G - \Delta G_{\text{ref}} > 0.5, \quad (3)$$

$$\text{Score} = 0.5 \text{ if } |\Delta G - \Delta G_{\text{ref}}| \leq 0.5, \quad (4)$$

$$\text{Score} = 1.0 \text{ if } \Delta G - \Delta G_{\text{ref}} < -0.5, \quad (5)$$

where ΔG_{ref} is the docking score of the protein with co-crystallized active reference ligand, and ΔG is the docking score for a ligand bound to that protein binding site. Shades were then set to white, grey, and black.

4.6. Biological Evaluation

4.6.1. Cultures for Parasites and Cells

Leishmania donovani promastigotes (MHOM/ET/67/HU3/LV9) were cultured with 5% CO₂ in M199 complete medium containing M199 medium supplemented with 100 μ M adenosine, 0.5 mg/L hemin, 40 mM Hepes pH 7.4 and 10% heat inactivated fetal bovine serum (HIFBS) in the dark at 26 °C. Late log promastigotes diluted at 1×10^6 /mL in M199 complete medium acidified at pH 5.5 and cultured at 37 °C with 5% CO₂ provided the cultures of axenic amastigotes of *L. donovani*.

4.6.2. Macrophage

RAW 264.7 macrophages were grown with 5% CO₂ in DMEM complete medium containing Dulbecco's Modified Eagle's Medium (DMEM) with 100 U/mL penicillin-streptomycin, and 10% HIFBS at 37 °C.

4.6.3. In Vitro Anti-Leishmanial Compound Testing on Axenic and Intramacrophage Amastigotes:

Previously described protocols [35] were adapted for evaluation of compounds on *L. donovani*. For axenic amastigotes, two-fold serial dilutions of the compounds were done in 96-well microplates with 100 μ L of complete medium (see above). Axenic amastigotes were then added to each well at a density of 106 mL in a 200 μ L final volume. After 72 h of incubation with 5% CO₂ at 37 °C, 20 μ L of resazurin (450 μ M) was added to each well and further incubated with 5% CO₂ at 37 °C for 24 h in the dark. Resazurin is reduced to resorufin in living cells and can be then monitored by measuring

OD570nm (resorufin) and OD600nm (resazurin; Lab systems Multiskan MS). Compound activity was expressed as IC₅₀ (μM). The drug used for reference was Amphotericin B (AmB).

For intramacrophage amastigote testing, RAW 264.7 macrophages were incubated with 5% CO₂ at 37 °C for 24 h after being plated in 96-well microplates at a density of 2 × 10⁴ cells per well. Axenic amastigotes were differentiated as described above, centrifuged at 2000 g for 10 min, re-suspended in DMEM complete medium, and added to each well to reach a 16:1 parasite to macrophage ratio. 24 h of infection with 5% CO₂ at 37 °C proceeded, and extracellular parasites were removed. This was followed by addition to each well of DMEM complete medium (100 μL) containing two-fold serial dilutions of the compounds from a maximal concentration of 100 μM. The medium was removed following treatment of 48 h and replaced by Direct PCR Lysis Reagent (100 μL; Euromedex) before 3 freeze-thaw cycles at room temperature, addition of 50 μg/mL proteinase K, and a final incubation at 55 °C overnight to allow cell lysis. 10 μL of each cell extract was then added to 40 μL of Direct PCR Lysis reagent containing Sybr Green I (0.05%; Invitrogen). Mastercycler[®] realplex (Eppendorf) was used to monitor DNA fluorescence. Compound activity was expressed as IC₅₀ (μM). The drug used for reference was Amphotericin B (AmB).

4.6.4. Cytotoxicity Tests

RAW 264.7 macrophages were used for cytotoxicity testing. Cells were plated in 96-well microplates at a density of 2 × 10⁴ cells per well. Twenty-four hours of incubation with 5% CO₂ at 37 °C followed, and then the medium was removed and to each well, 100 μL of DMEM complete medium containing two-fold serial dilutions of the compounds were added. This was followed by 48 h of incubation with 5% CO₂ at 37 °C, after which 10 μL of resazurin (450 μM) was added to each well, and further incubated with 5% CO₂, in the dark for 4 h at 37 °C. Cell viability was then monitored as above. Compound cytotoxicity was expressed as CC₅₀ (Cytotoxic Concentration 50%: concentration inhibiting the macrophages growth by 50%).

5. Conclusions

There is a growing and urgent need for new chemical compounds for the treatment of several leishmaniases given the lack of vaccines and the side-effects and lack of efficacy of presently-used chemicals. Four different series of compounds were synthesized and tested as potent anti-leishmanial compounds. While the indolizine compound does not present good activity, and the aldoxime and 1,2,4-oxadiazole tested are both equally moderately active, *N*-acylhydrazones presented much better activities. Among them, compound 2 is better than the others and could be considered as an interesting molecular template for further development of new anti-leishmanial agents due to its observed experimental activity against parasite *Leishmania donovani* axenic and intramacrophage amastigotes, better selectivity index, and predicted interactions with targets and anti-targets. It has a selectivity index of 2.2. Further work may be carried out to improve the toxicity and potency profiles of these compounds.

Author Contributions: M.B., A.T.G.-S., and S.G. conceived and designed the study; A.T.G.-S., M.S., M.D., C.M., R.B., S.C., S.S., and A.B. performed experiments; A.T.G.-S., P.M.L., M.B., M.S., K.N., S.G., and S.S. analyzed the data; A.T.G.-S., M.B., P.M.L., S.G., and M.S. wrote the paper; all authors read and approved the final manuscript.

Funding: This research was funded by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant no. 173001) and the Haridus- ja Teadusministeerium (Grant IUT34-14).

Acknowledgments: The authors would like to gratefully acknowledge the EU COST Action CM1307 “Targeted chemotherapy towards diseases caused by endoparasites” for traveling and publishing costs in open access. The NMR Facility (ICT, Toulouse, France, <http://ict.ups-tlse.fr/?RMN-103>) is gratefully acknowledged for carrying out the NMR analysis.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. WHO, Leishmaniasis: Situation and Trends. World Health Organization, 2017. Available online: https://www.who.int/gho/neglected_diseases/leishmaniasis/en/ (accessed on 19 February 2019).
2. WHO Fact Sheet. 2018. Available online: <http://www.who.int/en/news-room/fact-sheets/detail/leishmaniasis> (accessed on 19 February 2019).
3. Chappuis, F.; Sundar, S.; Hailu, A.; Ghalib, H.; Rijal, S.; Peeling, R.W.; Alvar, J.; Boelaert, M. Visceral leishmaniasis: What are the needs for diagnosis, treatment and control? *Nat. Rev. Microbiol.* **2007**, *5*, 873–882. [CrossRef] [PubMed]
4. Machado-Silva, A.; Guimarães, P.P.; Tavares, C.A.; Sinisterra, R.D. New perspectives for leishmaniasis chemotherapy over current anti-leishmanial drugs: A patent landscape. *Expert Opin. Ther. Pat.* **2015**, *25*, 247–260. [CrossRef] [PubMed]
5. Rajasekaran, R.; Chen, Y.P. Potential therapeutic targets and the role of technology in developing novel antileishmanial drugs. *Drug Discov. Today* **2015**, *20*, 958–968. [CrossRef] [PubMed]
6. De Menezes, J.; Guedes, C.; Petersen, A.; Fraga, D.; Veras, P. Advances in development of new treatment for Leishmaniasis. *Biomed. Res. Int.* **2015**, *2015*, 815023. [CrossRef] [PubMed]
7. Ponte-Sucre, A.; Gamarro, F.; Dujardin, J.C.; Barrett, M.P.; López-Vélez, R.; García-Hernández, R.; Pountain, A.W.; Mwenechanya, R.; Papadopolou, B. Drug resistance and treatment failure in leishmaniasis: A 21st century challenge. *PLoS Negl. Trop. Dis.* **2017**, *14*, e0006052. [CrossRef]
8. Rahman, R.; Goyal, V.; Haque, R.; Jamil, K.; Faiz, A.; Samad, R.; Ellis, S.; Balasegaram, M.; Boer, M.D.; Rijal, S.; et al. Safety and efficacy of short course combination regimens with AmBisome, miltefosine and paromomycin for the treatment of visceral leishmaniasis (VL) in Bangladesh. *PLoS Negl. Trop. Dis.* **2017**, *11*, e0005635. [CrossRef]
9. Rogers, M.; Kropf, P.; Choi, B.S.; Dillon, R.; Podinovskaia, M.; Bates, P.; Müller, I. Proteophosphoglycans regurgitated by Leishmania-infected sand flies target the L-arginine metabolism of host macrophages to promote parasite survival. *PLoS Pathog.* **2009**, *5*, e1000555. [CrossRef]
10. Ilari, A.; Fiorillo, A.; Genovese, I.; Colotti, G. Polyamine-trypanothione pathway: An update. *Future Med. Chem.* **2017**, *9*, 61–77. [CrossRef]
11. Da Silva, E.R.; Castilho, T.M.; Pioker, F.C.; Tomich de Paula Silva, C.H.; Floeter-Winter, L.M. Genomic organisation and transcription characterisation of the gene encoding *Leishmania (Leishmania) amazonensis* arginase and its protein structure prediction. *Int. J. Parasitol.* **2002**, *32*, 727–737. [CrossRef]
12. Bocedi, A.; Dawood, K.F.; Fabrini, R.; Federici, G.; Gradoni, L.; Pedersen, J.Z.; Ricci, G. Trypanothione efficiently intercepts nitric oxide as a harmless iron complex in trypanosomatid parasites. *FASEB J.* **2010**, *24*, 1035–1042. [CrossRef]
13. Colotti, G.; Ilari, A. Polyamine metabolism in *Leishmania*: From arginine to trypanothione. *Amino Acids* **2011**, *40*, 269–285. [CrossRef]
14. Reguera, R.M.; Balaña-Fouce, R.; Showalter, M.; Hickerson, S.; Beverley, S.M. *Leishmania major* lacking arginase (ARG) are auxotrophic for polyamines but retain infectivity to susceptible BALB/c mice. *Mol. Biochem. Parasitol.* **2009**, *165*, 48–56. [CrossRef]
15. Chawla, B.; Madhubala, R. Drug targets in *Leishmania*. *J. Parasit. Dis.* **2010**, *34*, 1–13. [CrossRef] [PubMed]
16. Lepesheva, G.I.; Hargrove, T.Y.; Rachakonda, G.; Wawrzak, Z.; Pomel, S.; Cojean, S.; Nde, P.N.; Nes, W.D.; Locuson, C.W.; Calcutt, M.W.; et al. VFV as a New Effective CYP51 Structure-Derived Drug Candidate for Chagas Disease and Visceral Leishmaniasis. *J. Infect. Dis.* **2015**, *212*, 1439–1448. [CrossRef] [PubMed]
17. Cottrell, D.M.; Capers, J.; Salem, M.M.; DeLuca-Fradley, K.; Croft, S.L.; Werbovetz, K.A. Antikinetoplastid activity of 3-aryl-5-thiocyanatomethyl-1,2,4-oxadiazoles. *Bioorg. Med. Chem.* **2004**, *12*, 2815–2824. [CrossRef] [PubMed]
18. Sharma, V.; Kumar, V. Indolizine: A biologically active moiety. *Med. Chem. Res.* **2014**, *23*, 3593–3606. [CrossRef]
19. Jaisankar, P.; Pal, B.; Manna, R.K.; Pradhan, P.K.; Medda, S.; Basu, M.K.; Giri, V.S. Synthesis of antileishmanial(5R)-(-)-5-carbomethoxy-3-formyl-5,6-dihydroindolo-[2,3-a]-indolizine. *ARKIVOC* **2003**, *ix*, 150–157.
20. Glisic, S.; Sencanski, M.; Perovic, V.; Stevanovic, S.; García-Sosa, A.T. Arginase Flavonoid Anti-Leishmanial in Silico Inhibitors Flagged against Anti-Targets. *Molecules* **2016**, *21*, 589. [CrossRef] [PubMed]

21. Sreekanth Thota, S.; Rodrigues, D.A.; Murteira Pinheiro, P.S.; Lima, L.M.; Fraga, C.A.M.; Barreiro, E.J. N-Acylhydrazones as drugs. *Bioorg. Med. Chem. Lett.* **2018**, *28*, 2797–2806. [[CrossRef](#)]
22. Andrade, M.M.; Barros, M.T. Fast synthesis of N-acylhydrazones employing a microwave assisted neat protocol. *J. Comb. Chem.* **2010**, *12*, 245–247. [[CrossRef](#)] [[PubMed](#)]
23. Oliveira, P.F.M.; Guidetti, B.; Chamayou, A.; André-Barrès, C.; Madacki, J.; Korduláková, J.; Mori, G.; Orena, B.S.; Chiarelli, L.R.; Pasca, M.R.; et al. Mechanochemical synthesis and biological C. evaluation of novel isoniazid derivatives with potent antitubercular activity. *Molecules* **2017**, *22*, 1457. [[CrossRef](#)]
24. Bora, R.O.; Dar, B.; Pradhan, V.; Farooqui, M. 1,2,4-Oxadiazoles: Synthesis and Biological applications. *Mini-Rev. Med. Chem.* **2014**, *14*, 355–369. [[CrossRef](#)] [[PubMed](#)]
25. Barros, C.J.P.; Rufino de Freitas, J.J.; De Oliveira, R.N.; De Freitas Filho, J.R. Synthesis of amidoximes using an efficient and rapid ultrasound method. *J. Chil. Chem. Soc.* **2011**, *56*, 721–722. [[CrossRef](#)]
26. Charlson, A.J.; Harington, J.S. The anti-inflammatory and analgesic activity of some benzimidazoles, and their ability to protect erythrocytes from hemolysis by silica powder. *Carbohydr. Res.* **1975**, *43*, 383–387. [[CrossRef](#)]
27. Alcalde, E.; Pérez-García, L.; Miravittles, C.; Rius, J.; Valentí, E. Heterocyclic Betaines. 13. Synthesis, Electronic and Molecular Structures of Methylenepyridinium and Methylenimidazolium Azolate Inner Salts. *J. Organ. Chem.* **1992**, *57*, 4829–4834. [[CrossRef](#)]
28. Daina, A.; Michielin, O.; Zoete, V. SwissADME: A free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* **2017**, *7*, 42717. [[CrossRef](#)]
29. Bolognesi, M.L.; Lizzi, F.; Perozzo, R.; Brun, R.; Cavalli, A. Synthesis of a small library of 2-phenoxy-1,4-naphthoquinone and 2-phenoxy-1,4-anthraquinone derivatives bearing anti-trypanosomal and anti-leishmanial activity. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 2272–2276. [[CrossRef](#)] [[PubMed](#)]
30. Veljkovic, V. *A Theoretical Approach to Preselection of Carcinogens and Chemical Carcinogenesis*; Gordon & Breach: New York, NY, USA, 1980.
31. Veljkovic, V.; Slavic, I. Simple general-model pseudopotential. *Phys. Rev. Lett.* **1972**, *29*, 105–107. [[CrossRef](#)]
32. *Virtual Screening Workflow*; Schrödinger, LLC: New York, NY, USA, 2018.
33. García-Sosa, A.T.; Sild, S.; Maran, U. Docking and virtual screening using distributed grid technology. *QSAR Comb. Sci.* **2009**, *28*, 815–821. [[CrossRef](#)]
34. García-Sosa, A.T.; Maran, U. Improving the use of ranking in virtual screening against HIV-1 integrase with triangular numbers and including ligand profiling with Antitargets. *J. Chem. Inf. Model.* **2014**, *54*, 3172–3185. [[CrossRef](#)] [[PubMed](#)]
35. Balaraman, K.; Vieira, N.C.; Moussa, F.; Vacus, J.; Cojean, S.; Pomel, S.; Bories, C.; Figadère, B.; Kesavan, V.; Loiseau, P.M. In vitro and in vivo antileishmanial properties of a 2-n-propylquinoline hydroxypropyl β -cyclodextrin formulation and pharmacokinetics via intravenous route. *Biomed. Pharmacother.* **2015**, *76*, 127–133. [[CrossRef](#)] [[PubMed](#)]

Sample Availability: Samples of the compounds are available from the authors.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).